



GLOBAL JOURNAL OF HUMAN SOCIAL SCIENCE
GEOGRAPHY, GEO-SCIENCES, ENVIRONMENTAL DISASTER
MANAGEMENT

Volume 13 Issue 7 Version 1.0 Year 2013

Type: Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-460X & Print ISSN: 0975-587X

The Utilization of MODIS and Landsat TM/ETM+ for Cotton Fractional Yield Estimation in Burewala

By Farooq Ahmad, Qurat-ul-ain Fatima, Hira Jannat Butt, Kashif Shafique,
Sajid Rashid Ahmad, Shafeeq-Ur-Rehman, Rao Mansor Ali Khan & Abdul Raoof

University of the Punjab, New Campus, Lahore, Pakistan

Abstract- Estimates of crop yield are desirable for managing agricultural lands. Remote sensing is the one technology that can give an unbiased view of large areas, with spatially explicit information distribution and time repetition, and has thus been widely used to estimate crop yield and offers great potential for monitoring production, yet the uncertainties associated with large-scale crop yield estimates are rarely addressed. In this study, we tried to estimate cotton cropped area using the supervised classification; planting dates for 11 years (1998 to 2009) of Landsat imagery, and fractional yield using MODIS (Terra) the normalized difference vegetation index (NDVI), and enhanced vegetation index (EVI) in an intensive agricultural region of Burewala, Punjab province of Pakistan. Vegetation indices are widely used for assessing and monitoring ecological variables such as vegetation cover and above-ground biomass. Monitoring the spatial distribution of cotton yield helps identifying sites with yield constraints. The newly available satellite images from the MODIS sensor provide enhanced atmospheric correction, cloud detection, improved geo-referencing, comprehensive data quality control and the enhanced ability to monitor vegetation development. The high temporal resolution of the MODIS datasets can provide an efficient and consistent way for biomass and fractional yield monitoring and assessment. The reflected radiation provides an indication of the type and density of canopy. The condition, distribution, structure and the development of the vegetation through the phenological stages can affect the relation between yield and NDVI. The high spatial resolution Landsat images were applied to extract the area under cotton cultivation within the landscape and to determine the cotton fraction among other land uses within the coarse spatial resolution MODIS pixels.

Keywords: Crop yield, EVI, cotton fraction, Landsat, MODIS dataset, NDVI, remote sensing.

GJHSS-B Classification : FOR Code: 059999



THE UTILIZATION OF MODIS AND LANDSAT TM/ETM+ FOR COTTON FRACTIONAL YIELD ESTIMATION IN BUREWALA

Strictly as per the compliance and regulations of:



The Utilization of MODIS and Landsat TM/ETM+ for Cotton Fractional Yield Estimation in Burewala

Farooq Ahmad ^α, Qurat-ul-ain Fatima ^σ, Hira Jannat Butt ^ρ, Kashif Shafique ^ω, Sajid Rashid Ahmad [¥], Shafeeq-Ur-Rehman [§], Rao Mansoor Ali Khan ^x & Abdul Raouf ^v

Abstract- Estimates of crop yield are desirable for managing agricultural lands. Remote sensing is the one technology that can give an unbiased view of large areas, with spatially explicit information distribution and time repetition, and has thus been widely used to estimate crop yield and offers great potential for monitoring production, yet the uncertainties associated with large-scale crop yield estimates are rarely addressed. In this study, we tried to estimate cotton cropped area using the supervised classification; planting dates for 11 years (1998 to 2009) of Landsat imagery, and fractional yield using MODIS (Terra) the normalized difference vegetation index (NDVI), and enhanced vegetation index (EVI) in an intensive agricultural region of Burewala, Punjab province of Pakistan. Vegetation indices are widely used for assessing and monitoring ecological variables such as vegetation cover and above-ground biomass. Monitoring the spatial distribution of cotton yield helps identifying sites with yield constraints. The newly available satellite images from the MODIS sensor provide enhanced atmospheric correction, cloud detection, improved geo-referencing, comprehensive data quality control and the enhanced ability to monitor vegetation development. The high temporal resolution of the MODIS datasets can provide an efficient and consistent way for biomass and fractional yield monitoring and assessment. The reflected radiation provides an indication of the type and density of canopy. The condition, distribution, structure and the development of the vegetation through the phenological stages can affect the relation between yield and NDVI. The high spatial resolution Landsat images were applied to extract the area under cotton cultivation within the landscape and to determine the cotton fraction among other land uses within the coarse spatial resolution MODIS pixels.

Keywords: Crop yield, EVI, cotton fraction, Landsat, MODIS dataset, NDVI, remote sensing.

1. INTRODUCTION

The high temporal resolution of the MODIS datasets can provide an efficient and consistent way for biomass and yield monitoring (Dalezios et al., 2001; Alexandridis and Chemin, 2002). Increased knowledge about the spatial distribution of cotton yield

(Shi et al., 2007) in Burewala, Punjab province of Pakistan, supports the optimal provision of resources (Macdonald and Hall, 1980; Hutchinson, 1991; Lobell et al., 2003; Hongo and Niwa, 2012).

Remote sensing imagery offers unique possibilities for spatial and temporal characterization of the changes. The basic requirement is the availability of different dates of imagery which permit continuous monitoring of change and environmental developments over time (Lu et al., 2004; Nasr and Helmy, 2009; Ahmad, 2012d). Long-term observations of remotely sensed vegetation dynamics have held an increasingly prominent role in the study of terrestrial ecology (Budde et al., 2004; Prasad et al., 2007; Ouyang et al., 2012; Ahmad, 2012). A major limitation of such studies is the limited availability of sufficiently consistent data derived from long-term remote sensing (Ouyang et al., 2012; Ahmad, 2012). The benefit obtained from a remote sensing sensor, thereby, largely depends on its spectral resolution (Jensen, 2005; Ahmad, 2012), which determines the sensor's capability to resolve spectral features of land surfaces (Fontana, 2009; Ahmad, 2012). One of the key factors in assessing vegetation dynamics and its response to climate change is the ability to make frequent and consistent observations (Thomas and Leason, 2005; Ouyang et al., 2012; Ahmad, 2012). Photosynthetically active vegetation is characterized by very low reflectance values (Jensen, 2005; Ahmad, 2012a) in the red part of the electromagnetic spectrum due to the absorption of solar radiation by the leaf pigments involved in photosynthesis (Lillesand et al., 2004; Ahmad, 2012a), and by increased reflectance in the near infrared portion of the spectrum due to reflection of incoming solar radiation at the leaf internal structures (Gitelson and Merzlyak, 1996; Fontana, 2009; Ahmad, 2012a). With regard to the monitoring of terrestrial vegetation, the Normalized Difference Vegetation Index (Rouse et al., 1973; Tucker, 1979; Ahmad, 2012a) is the most commonly used index and serves as a measure of photosynthetic activity within a certain area (Fontana, 2009; Ahmad, 2012a).

Los et al. (1994) and Sellers et al. (1996) were the first to derive land surface parameters with realistic seasonal and spatial variations for the globe from NDVI data collected by the AVHRR satellite (Los et al., 2000;

Author α: Department of Geography, University of the Punjab, New Campus, Lahore, Pakistan. e-mail: drylandpk@yahoo.com

Author σ ρ: GIS Centre, PUCIT, University of the Punjab, Lahore, Pakistan.

Author ω: Forman Christian College (A Chartered University), Lahore, Pakistan.

Authors ¥ § x v: Institute of Geology, University of the Punjab, New Campus, Lahore, Pakistan.

Ahmad, 2012a). Estimation of land surface vegetation parameters from satellite is based on the spectral properties of vegetation; vegetation strongly absorbs visible light, using the energy for photosynthesis, and strongly reflects near-infrared (NIR) radiation (Rouse et al., 1973; Los, 1998; Los et al., 2000; Ahmad, 2012a; Ahmad, 2012b). Changes in vegetation spectral response caused by phenology can conceal the longer term changes in the landscape (Hobbs, 1989; Lambin, 1996; Dennison and Roberts, 2003; Ahmad, 2012d). Multi-temporal data that captures these spectral differences can improve reparability of vegetation types over classifications based on single date imagery (DeFries et al., 1995; Ahmad, 2012d).

The Moderate Resolution Imaging Spectroradiometer (MODIS) is a key instrument onboard the Terra satellite platform (Huete et al., 2006; Carrão et al., 2008; Ahmad, 2012c). Terra's orbit around the Earth is timed so that it passes from north to south across the equator in the morning (Salomonson and Toll, 1991; GSFC/NASA, 2003; Huete, 2005; Ahmad, 2012c). There are now over 13 years of MODIS Terra data (first image, February 24, 2000), available producing high quality scientific products with calibration specifications of 2% reflectance and 5% radiance and geolocation of 50 m (Huete, 2005; Ahmad, 2012c).

The MODIS sensor has 36 spectral bands extending from the visible to the thermal infrared wavelengths [between 0.405 and 14.385 μm] (Running et al., 1994; Zhan et al., 2000). The MODIS land bands have a heritage related to the Landsat TM, with additional spectral capabilities added in the short-wave and long-wave infrared (Justice et al., 1998; Jin and Sader, 2005). Several researchers have used MODIS to detect changes in land cover (Zhan et al., 2000; 2002; Roy et al., 2002; Korontzi et al., 2004; Jin and Sader, 2005).

A remote sensing sensor is a key device that captures data about an object or scene remotely. Since objects have their unique spectral features, they can be identified from remote sensing imagery according to their unique spectral characteristics (Xie, 2008; Ahmad, 2013). A good case in vegetation mapping using remote sensing technology is the spectral radiances in the red and near-infrared (NIR) regions, in addition to others. The radiances in these regions could be incorporated into the spectral vegetation indices (VI) that are directly related to the intercepted fraction of photosynthetically active radiation (Asrar et al., 1984; Galio et al., 1985; Xie, 2008; Ahmad, 2013). Such evaluations often require the use of vegetation indices calculated from archived satellite data.

The NDVI is chlorophyll sensitive; the EVI (Liu and Huete, 1995; Justice et al., 1998; Huete et al., 1999) is more responsive to canopy structural variations, including canopy type, plant physiognomy and canopy architecture (Gao et al., 2000; Huete et al., 2002). The

two VIs complement each other in global vegetation studies and improve upon the detection of vegetation changes and extraction of canopy biophysical parameters (Huete et al., 1999; 2002).

The NDVI can be a useful tool to couple climate and vegetation distribution and performance at large spatial and temporal scales (Pettorelli et al., 2005; Aguilar et al., 2012; Ahmad, 2012; Ahmad, 2013a) because vegetation vigor and productivity are related to temperature-precipitation and evapotranspiration. The NDVI serves as a surrogate measure of these factors at the landscape scale (Wang et al., 2003; Groeneveld and Baugh, 2007; Aguilar et al., 2012; Ahmad, 2012; Ahmad, 2013a). The NDVI product works optimally with cloud filtering, radiometric calibration, precise geolocation, and a snow mask. In addition, the product performs best using top-of-canopy reflectance inputs, corrected for atmospheric ozone, molecular scattering, aerosol, and water vapour (Huete et al., 2006; Ahmad, 2012; Ahmad, 2013a).

Accurate evaluation of vegetation response across multiple-year time scales is crucial for analyses of global change (Running and Nemani, 1991; Sellers et al., 1994; Stow, 1995; Justice et al., 1998; Fensholt, 2004; Baugh and Groeneveld, 2006; Ahmad, 2012e), effects of human activities (Moran et al., 1997; Milich and Weiss 2000; Thiam, 2003; Baugh and Groeneveld, 2006; Ahmad, 2012e) and ecological relationships (Baret and Guyot, 1991; Asrar et al., 1992; Begue, 1993; Epiphanyo and Huete, 1995; Gillies et al., 1997; Baugh and Groeneveld, 2006; Ahmad, 2012e).

a) Study Area

Burewala (Figure 1) lies in the Punjab province of Pakistan from 29° 52' 28" to 30° 22' 12" North latitude and 72° 30' 04" to 72° 59' 35" East longitude.




Figure 1: Burewala - Landsat TM 2nd August, 2009 image

Source: <http://glovis.usgs.gov/>

II. RESEARCH DESIGN AND METHODS

In this research paper four Landsat TM/ETM+ scenes of 20th August, 1998; 15th August, 1999; 20th August, 2001 and 2nd August, 2009 (path 150; row 39) were used to detect and recognize the cotton-pixels and cotton cropped areas in Burewala. These vital steps are: image registration and image enhancement as discussed by Macleod and Congalton (1998), Mahmoodzadeh (2007) and Al-Awadhi et al., 2011. These scenes were corrected and geo-referenced using projection UTM, zone 43 and datum WGS 84. The Landsat has a long history of dataset, it is very helpful to map long-term vegetation cover and study the spatio-temporal vegetation changes (Schroeder et al., 2006; Xie, 2008; Ahmad, 2012d).

Radiometric enhancement is performed to produce a homogenous radiometric set of data (Richard et al., 2005; Ahmad, 2012d). So that, false changes are not introduced by factors such as: modification of the spectral distribution due to atmospheric conditions, different path radiance and seen angle variation (Ulbricht and Heckendorf, 1998; Ahmad, 2012d). Ideally, all images would be calibrated to standard reflectance units (Nasr and Helmy, 2009; Ahmad, 2012d). However, when comparing images to detect change, it is sufficient to convert raw digital counts to be consistent with a chosen reference image (Symeonakis et al., 2006; Ahmad, 2012d). Histograms equalization and histogram matching were generated in order to identify the suitable contrast stretching level to optimize the balance (Singh, 1989; Ahmad, 2012d) for Landsat TM/ETM+ images. Landsat TM/ETM+ different bands have used in order to estimate the vegetation quantities parameter based on

vegetation indices (Ahmad, 2012e). While all dimensions of remotely sensed data are relevant, for practical purposes it is the temporal information that has been most useful for monitoring of major crop types with remote sensing (Smith and Ramey, 1982; Badhwar, 1984; Hall and Badhwar, 1987; Price et al., 1997; Wardlow et al., 2007; Singh, 2012).

The MODIS (Terra) NDVI/EVI (MOD13A1) data products for research area were acquired, in this case data were downloaded from the Land Processes Distributed Active Archive Center (LPDAAC). Tile number covering this area is h24v05, reprojected from the Integerized Sinusoidal projection to a Geographic Lat/Lon projection, and Datum WGS84 (GSFC/NASA, 2003; Ahmad, 2012; 2012d). A gapless time series of MODIS (Terra) NDVI/EVI composite raster data from February, 2000 to February, 2013 with a spatial resolution of 500 m (Table 1) was utilized for calculation of the cotton fractional yield (Figure 2). The datasets provide frequent information at the spatial scale at which the majority of human-driven land cover changes occur (Townshend and Justice, 1988; Verbesselt et al., 2010; Ahmad, 2012). MODIS products are designed to provide consistent spatial and temporal comparisons between different global vegetation conditions that can be used to monitor photosynthetic activity and forecast crop yields (Vazifiedoust et al., 2009; Cheng and Wu, 2011). Details documenting the MODIS (Terra) NDVI/EVI compositing process and Quality Assessment Science Data Sets can be found at NASA's MODIS web site (MODIS, 1999; USGS, 2008).

Table 1: MODIS (Terra) Bands used in this research study

Bandwidth specifications (nm)	Band 3: 459-479
	Band 4: 545-565
	Band 5: 1230-1250
	Band 6: 1628-1652
	Band 7: 2105-2155
Spatial resolution (m)	500
Radiometric resolution (bits)	12
Time window	16-days

Earth location data is available at sub-pixel accuracy. This extraordinary geolocation accuracy is achieved due to several reasons (Wolfe et al., 1995; 2002; Fontana, 2009; Ahmad, 2012): First, the spacecrafts carrying MODIS are very stable and provide highly precise external orientation knowledge. Second, the MODIS instrument was designed to give precise interior orientation knowledge (Khlopenkov and Trishchenko, 2008; Fontana, 2009; Ahmad, 2012). Third, an accurate global DEM (Logan, 1999; Fontana, 2009; Ahmad, 2012) is used to model and remove relief-induced distortions. Fourth, a global set of GCPs based

on Landsat imagery served to determine biases in the sensor orientation, which were finally used to improve geolocation processing (Ackerman et al., 1998; 2006; Fontana, 2009; Ahmad, 2012).

The NDVI is successful as a vegetation measure is that it is sufficiently stable to permit meaningful comparisons of seasonal and inter-annual changes in vegetation growth and activity (Choudhury, 1987; Jakubauskas et al., 2002; Chen et al., 2006; Zoran and Stefan, 2006; Nicandrou, 2010; Ahmad, 2012; Ahmad, 2012d; 2012e). The strength of the NDVI is in its ratioing concept (Moran et al., 1992; Ahmad, 2012),

which reduces many forms of multiplicative noise (illumination differences, cloud shadows, atmospheric attenuation, and certain topographic variations) present in multiple bands (Chen et al., 2002; Nicandrou, 2010; Ahmad, 2012; Ahmad, 2012d).




Figure 2 : Scheme for research design and methods

The NDVI, spectral vegetation index which measures soil and vegetation moisture (Singh, 1989; Lyon et al., 1998; Mambo and Archer, 2007; Ahmad, 2013a), has been widely used for environmental change monitoring (Young, 1998; Lillesand and Kiefer, 2000; Eastman, 2003; Lillesand et al., 2004; Mambo and Archer, 2007; Ahmad, 2013a). The index can be used to identify areas showing distressed or degraded vegetation, leading to identification of possible degraded areas (Barrow, 1991; Booth et al., 1994; Mambo and Archer, 2007; Ahmad, 2013a). The NDVI captures the marked contrast between the strong absorptance in the visible wavelengths and strong reflectance in the near-infrared wavelengths which uniquely characterize the presence of photosynthetically

active vegetation (Tucker, 1979; Wessels et al., 2004; Ahmad, 2013a). The NDVI is an indicator of vegetation health, because degradation of ecosystem vegetation, or a decrease in green, would be reflected in a decrease in NDVI value (Meneses-Tovar, 2011; Ahmad, 2013a).

The NDVI is highly correlated with vegetation parameters such as green leaf biomass and green leaf area (Justice et al., 1985; Ahmad, 2013a), and it also is directly related to plant vigor, density, and growth conditions (Holben, 1986; Ahmad, 2013a), now it is widely accepted as a primary tool for monitoring land degradation (Huang et al., 2010; Ahmad, 2013a). In arid and semi-arid lands, seasonal sums of multi-temporal NDVI are strongly correlated with vegetation production (Prince and Tucker, 1986; Prince, 1991; Nicholson and Farrar, 1994; Nicholson et al., 1998; Wessels et al., 2004; Ahmad, 2013a).

The NDVI values range from -1 to +1; because of high reflectance in the NIR portion of the EMS, healthy vegetation is represented by high NDVI values between 0.1 and 1 (Liu and Huete, 1995; USGS, 2008; 2010; Ahmad, 2012). Conversely, non-vegetated surfaces such as water bodies yield negative values of NDVI because of the electromagnetic absorption property of water. Bare soil areas represent NDVI values which are closest to 0 due to high reflectance in both the visible and NIR portions of the EMS (Townshend, 1992; Ahmad, 2012). The NDVI is related to the absorption of photosynthetically active radiation and basically measures the photosynthetic capability of leaves, which is related to vegetative canopy resistance and water vapour transfer (Wan, 2003; Rahman et al., 2004; Ahmad, 2012; 2012d).

The NDVI is the most commonly used index of greenness derived from multispectral remote sensing data (USGS, 2010; Ahmad, 2013a), and is used in several studies on vegetation, since it has been proven to be positively correlated with density of green matter (Townshend et al., 1991; Huete et al., 1997; Huete et al., 2002; Debien et al., 2010; Ahmad, 2012a; 2012d; Zaeen, 2012; Ahmad, 2013a). The NDVI provides useful information for detecting and interpreting vegetation land cover it has been widely used in remote sensing studies (Dorman and Sellers, 1989; Myneni and Asrar, 1994; Gao, 1996; Sesnie et al., 2008; Karaburun, 2010; Ahmad, 2012b; Ahmad, 2013a).

The EVI is an 'optimized index' designed to enhance the vegetation signal with improved sensitivity in high biomass regions and improved vegetation monitoring through a de-coupling of the canopy background signal and a reduction in atmosphere influences (Liu and Huete, 1995; Justice et al., 1998; Huete et al., 1999; Ahmad, 2012e), and minimizes canopy-soil and aerosol variations and improves sensitivity over dense vegetation conditions (Qi et al., 1994; Ahmad, 2012c). The two products more effectively characterize a global range of vegetation states and

processes, and improve upon the extraction of canopy biophysical parameters (Jiang et al., 2008; Ahmad, 2012c).

ERDAS imagine 2013 and ArcGIS 10 software were used for application of NDVI model to detect the cotton cropped area and calculation for Landsat TM/ETM+ images (path 150; row 39) of 20th August, 1998; 15th August, 1999; 20th August, 2001 and 2nd August, 2009 respectively. The supervised classification was applied upon the images for the estimation of the cotton cropped area. Further, comparative analysis was carried out upon supervised classification to detect the deviation in the cotton cropped area in Burewala. Calculation of the cotton fractional yield using MODIS (Terra) NDVI/EVI pixel values of the selected villages; Chak 44/KB, Chak 305/EB, Chak 487/EB and Chak 529/EB of Burewala was carried out and linear forecast trendline was plotted to identify the variations in the cotton fractional yield dataset of Chak 44/KB from February 2000 to February 2013. Standard multispectral image processing techniques were generally developed to classify multispectral images into broad categories of surface condition (Shippert, 2004; Ahmad, 2012f). ERDAS imagine (ERDAS Imagine, Inc., 2010) and ENVI tools, applicable to a variety of applications, distinguish and identify the unique resource information present in the scene and map them throughout the image (Research System, Inc., 2004).

III. RESULTS

Figure 3, 4, 5 and 6 (Table 2) shows classified NDVI for Landsat TM/ETM+ scenes of 20th August, 1998; 15th August, 1999; 20th August, 2001 and 2nd August, 2009 respectively. The NDVI model was applied upon the Landsat TM/ETM+ using ERDAS imagine 2013 software while ArcGIS 10 was used for NDVI calculation. Further, ArcGIS symbology tool was used to develop NDVI classes and recognize the cotton cropped areas in Burewala. The NDVI is an effective vegetation measure since it is sufficiently stable to permit meaningful comparisons between seasonal and interannual changes in vegetation growth and activity (Cheng and Wu, 2011). This is because it can reduce different forms of multiplicative noise present in multiple bands (Myneni et al., 1995; Cheng and Wu, 2011). The significance of NDVI index may vary according to habitat type (Pettorelli et al., 2005; Hamel et al., 2009; Ahmad, 2013a). Vegetation indices may employ simple ratios of any two single wavelength combinations. These ratios were found to be fairly effective in normalizing the effect of reflectance variation in soil background (Colwell, 1973; Cheng and Wu, 2011).

Remote sensing data provides systematically high-quality spatial and temporal information about land surface features, including behaviour of agricultural crops and cumulative environmental impacts on crop

growing conditions (Liu and Kogan, 2002). Remote sensing techniques have been employed to estimate various plant parameters (Wiegant et al., 1979; Price, 1995; Mirik et al., 2007; Cheng and Wu, 2011) and crop yield (Cheng et al., 2004; Cheng, 2006; Cheng and Wu, 2011). Remote sensing provides quantitative information on agricultural crops instantaneously and nondestructively (Clevers, 1988; Mirik et al., 2006; Cheng and Wu, 2011), and the spatial and temporal distributions of crop production offer valuable information for agricultural management and biogeochemical modeling efforts (Lobell et al., 2003; Sönmez and Sari, 2006; Cheng and Wu, 2011).




Figure 3 : Classified NDVI 1998, Burewala




Figure 4 : Classified NDVI 1999, Burewala




Figure 5 : Classified NDVI 2001, Burewala




Figure 6 : Classified NDVI 2009, Burewala

Table 2 : NDVI values of Landsat images

Image Acquisition Date	Maximum NDVI	Minimum NDVI	Mean NDVI	Standard Deviation
20 th August, 1998 (Landsat TM)	0.61	-0.19	0.23	0.13
15 th August, 1999 (Landsat ETM+)	0.51	-0.26	0.11	0.12
20 th August, 2001 (Landsat ETM+)	0.56	-0.24	0.21	0.14
2 nd August, 2009 (Landsat TM)	0.54	-0.15	0.20	0.11

The NDVI (Sellers, 1985) as an indicator of vivid green vegetation and as a descriptor of ecosystem functions has proved to be very valuable for assessing ecological responses to environmental changes (Pettorelli et al., 2005; Alcaraz-Segura et al., 2009;

Höpfner and Scherer, 2011; Ahmad, 2012a). The NDVI can be used not only for accurate description of vegetation classification and vegetation phenology (Tucker et al., 1982; Tarpley et al., 1984; Justice et al., 1985; Lloyd, 1990; Singh et al., 2003; Los et al., 2005;

Ahmad, 2013a) but also effective for monitoring rainfall and drought, estimating net primary production of vegetation, crop growth conditions and crop yields, detecting weather impacts and other events important for agriculture and ecology (Kogan, 1987; Dabrowska-Zielinska et al., 2002; Singh et al., 2003; Chris and Molly, 2006; Baldi et al., 2008; Glenn et al., 2008; Ahmad, 2012b; Ahmad, 2013a).

Figure 7, 8, 9 and 10 shows the supervised classification for Landsat TM/ETM+ scenes of 20th August, 1998; 15th August, 1999; 20th August, 2001 and 2nd August, 2009 respectively. The supervised classification was applied upon the Landsat TM/ETM+ using ERDAS imagine 2013 software to detect cotton cropped areas in Burewala. The approach supervised classification which is part of post classification comparison method or direct classification method. This approach is based on the natural groupings of the spectral properties of the pixels which are usually selected by the RS software without any influence from the users (Al-Awadhi et al., 2011).

Figure 7 shows the supervised classification for Landsat TM image of 20th August, 1998. The findings showed that the cotton fields were 591.44 km² (47.77%), roads and settlements 402.34 km² (32.50%), and other crops 206.06 km² (16.65%) while the fallow land was 38.13 km² (3.08%). The accuracy assessment is given in Table 3.

Figure 8 shows the supervised classification for Landsat ETM+ image of 15th August, 1999. The findings showed that the cotton fields were 528.38 km² (42.68%), roads and settlements 404.55 km² (32.68%), and other crops 257.13 km² (20.77%) while the fallow land was 47.91 km² (3.87%). The accuracy assessment is given in Table 3.

Soil compaction negatively affects crop growth characteristics (Lowery and Schuler, 1991; Kulkarni and Bajwa, 2005), yield (Johnson et al., 1990; Kulkarni and Bajwa, 2005), and root development and distribution (Taylor and Gardener, 1963; Unger and Kaspar, 1994; Kulkarni and Bajwa, 2005). However, bare soil reflectance may be affected by the impact of tillage practices and moisture content (Barnes et al., 1996; Kulkarni and Bajwa, 2005). The wavelengths detected as responsive to soil compaction were close to each other, they may had similar information about the vegetation vigor. In the red portion of spectrum the wavelengths ranged from 620 to 700 nm (Thenkabail et al., 2000; Kulkarni and Bajwa, 2005). In this region of visible spectrum, a mixed spectral signature was produced from cotton bolls and the yellow and brownish green leaves. Some exposure of cotton stems as leaves had fallen and more soil exposure caused the mixed signature (Rosenthal and Gerik, 1991; Kulkarni and Bajwa, 2005).

Figure 9 shows the supervised classification for Landsat ETM+ image of 20th August, 2001. The findings


showed that the cotton fields were 702.61 km² (56.76%), roads and settlements 319.58 km² (25.81%), and other crops 175.56 km² (14.18%) while the fallow land was 40.22 km² (3.25%). The accuracy assessment is given in Table 3.

Figure 10 shows the supervised classification for Landsat TM image of 2nd August, 2009. The findings showed that the cotton fields were 494.11 km² (39.91%), roads and settlements 509.13 km² (41.13%), and other crops 213.58 km² (17.25%) while the fallow land was 21.15 km² (1.71%). The accuracy assessment is given in Table 3.

Figure 11 shows the comparative analysis of the supervised classification for Landsat TM/ETM+ scenes of 20th August, 1998; 15th August, 1999; 20th August, 2001 and 2nd August, 2009 respectively. The findings showed that the area of the cotton fields decreased from 591.44 km² (47.77%) on 20th August, 1998 to 494.11 km² (39.91%) on 2nd August, 2009 (Table 3).

Remote sensing provides a viable source of data from which updated land-cover information can be extracted efficiently and cheaply in order to invent and monitor these changes effectively (Mas, 1999; Ahmad, 2013a).






Cotton Fields
 Roads and Settlements
 Other Crops
 Fallow Land

Source:
<http://glovis.usgs.gov/>
 Landsat TM
 Acquisition Date: 20th August, 1998


Figure 7 : Supervised Classification 1998, Burewala



Cotton Fields
 Roads and Settlements
 Other Crops
 Fallow Land

Source:
<http://glovis.usgs.gov/>
 Landsat ETM+
 Acquisition Date: 15th August, 1999


Figure 8 : Supervised Classification 1999, Burewala



Cotton Fields
 Roads and Settlements
 Other Crops
 Fallow Land

Source:
<http://glovis.usgs.gov/>
 Landsat ETM+
 Acquisition Date: 20th August, 2001

Figure 9 : Supervised Classification 2001, Burewala



Cotton Fields
 Roads and Settlements
 Other Crops
 Fallow Land

Source:
<http://glovis.usgs.gov/>
 Landsat TM
 Acquisition Date: 2nd August, 2009

Figure 10 : Supervised Classification 2009, Burewala

Table 3 : Supervised Classification of Landsat TM/ETM+ images

Image Acquisition Date	Classes	Area (km ²)	Area (%)	Accuracy Assessment (%)
20 th August, 1998 (Landsat TM)	Cotton Fields	591.44	47.77	86.13
	Roads and Settlements	402.34	32.50	89.62
	Other Crops	206.06	16.65	85.47
	Fallow Land	38.13	3.08	88.75
	SUM	1237.97	100	-
15 th August, 1999 (Landsat ETM+)	Cotton Fields	528.38	42.68	87.38
	Roads and Settlements	404.55	32.68	92.35
	Other Crops	257.13	20.77	92.15
	Fallow Land	47.91	3.87	85.44
	SUM	1237.97	100	-

20 th August, 2001 (Landsat ETM+)	Cotton Fields	702.61	56.76	85.47
	Roads and Settlements	319.58	25.81	85.77
	Other Crops	175.56	14.18	85.16
	Fallow Land	40.22	3.25	90.15
	SUM	1237.97	100	-
2 nd August, 2009 (Landsat TM)	Cotton Fields	494.11	39.91	92.18
	Roads and Settlements	509.13	41.13	89.34
	Other Crops	213.58	17.25	88.32
	Fallow Land	21.15	1.71	85.47
	SUM	1237.97	100	-




Figure 11 : Comparative Analysis of Supervised Classification of Landsat TM/ETM+ Images

Figure 12 shows NDVI/EVI phenological profile for Chak 44/KB; Figure 13 shows NDVI/EVI phenological profile for Chak 305/EB; Figure 14 shows NDVI/EVI phenological profile for Chak 487/EB and Figure 15 shows NDVI/EVI phenological profile for Chak 529/EB, Burewala, Punjab province, Pakistan. The NDVI/EVI pixel values were used to calculate fractional yield (Shinners and Binversie, 2007) from February 2000 to February 2013. The NDVI pixel values showed theoretical yield and EVI pixel values showed actual yield (Table 4). Further, linear forecast trendline was plotted upon the fractional yield dataset of Chak 44/KB to investigate the general trend during the entire period (Figure 16). The linear forecast trendline showed that fractional yield at Chak 44/KB was decreasing. The tendency of decreasing in the fractional yield from February 2000 to December 2004 was low and the tendency from January 2005 to February 2013 was high.

Vegetation phenology refers to the relationship between climate and periodic development of photosynthetic biomass (Ahl et al., 2006; Ahmad, 2012a). Satellite monitoring of vegetation phenology has often made use of a vegetation index such as NDVI because it is related to the amount of green leaf biomass (Lillesand and Kiefer, 2000; Beurs and Henebry, 2004; Ahl et al., 2006; Ahmad, 2012a). The potential solution to provide more frequent high resolution surface observations is to fuse Landsat observations with data from other remote sensing systems, such as MODIS (Singh, 2012). The MODIS

NDVI datasets provides unique opportunities for monitoring terrestrial vegetation conditions at regional and global scales (Yang et al., 1997; Piao et al., 2006; Ahmad, 2012), and has widely been used in research areas of net primary production (Potter et al., 1993; Paruelo et al., 1997; Piao et al., 2006; Ahmad, 2012), vegetation coverage (Tucker et al., 1991; Myneni et al., 1997; Los et al., 2001; Zhou et al., 2001; Piao et al., 2003; Piao et al., 2006; Ahmad, 2012), biomass (Myneni et al., 2001; Dong et al., 2003; Piao et al., 2006), and phenology (Reed et al., 1994; Moulin et al., 1997; Piao et al., 2006; Ahmad, 2012).




Figure 12 : NDVI/EVI phenological profile for Chak 44/KB

Processed by the author




Figure 13 : NDVI/EVI phenological profile for Chak 305/EB

Processed by the author




Figure 14 : NDVI/EVI phenological profile for Chak 487/EB

Processed by the author




Figure 15 : NDVI/EVI phenological profile for Chak 529/EB

Processed by the author

Table 4 : MODIS (Terra) EVI/NDVI and Fractional Yield dataset of Chak 44/KB

Image Acquisition (Month/Year)	EVI Pixel Value	NDVI Pixel Value	Fractional Yield (%)	Image Acquisition (Month/Year)	EVI Pixel Value	NDVI Pixel Value	Fractional Yield (%)
Feb. 2000	4320	6127	70.51	Feb. 2007	5409	6605	81.89
May 2000	1108	1664	66.59	May 2007	2055	2288	89.82
Aug. 2000	3826	5102	74.99	Aug. 2007	4122	5381	76.6
Nov. 2000	2104	3222	65.3	Nov. 2007	1845	2858	64.56
Feb. 2001	4419	6595	67.01	Feb. 2008	4498	4819	93.34
May 2001	1520	1771	85.83	May 2008	1987	2501	79.45
Aug. 2001	4051	5442	74.44	Aug. 2008	4733	5528	85.62
Nov. 2001	1647	2898	56.83	Nov. 2008	2267	2602	87.13
Feb. 2002	4360	6397	68.16	Feb. 2009	4049	6222	65.08
May 2002	1958	2110	92.8	May 2009	1665	1923	86.58
Aug. 2002	4305	4834	89.06	Aug. 2009	3923	5463	71.81
Nov. 2002	2076	3414	60.81	Nov. 2009	1709	2434	70.21
Feb. 2003	4447	4902	90.72	Feb. 2010	4404	6864	64.16
May 2003	1537	1879	81.8	May 2010	1559	2345	66.48
Aug. 2003	4623	5571	82.98	Aug. 2010	4327	6165	70.19
Nov. 2003	1615	2783	58.03	Nov. 2010	2554	4549	56.14
Feb. 2004	4986	7037	70.85	Feb. 2011	3778	6040	62.55
May 2004	1294	1795	72.09	May 2011	2164	2983	72.54
Aug. 2004	4820	5857	82.29	Aug. 2011	4740	7275	65.15
Nov. 2004	2148	2983	72.01	Nov. 2011	2679	5094	52.59
Feb. 2005	5892	5400	109.11	Feb. 2012	3102	5055	61.36
May 2005	1598	1994	80.14	May 2012	1785	2768	64.49
Aug. 2005	5599	5380	104.07	Aug. 2012	3564	5251	67.87
Nov. 2005	2242	3144	71.31	Nov. 2012	2639	4480	58.91
Feb. 2006	5230	6031	86.72	Feb. 2013	4059	6476	62.68
May 2006	1607	2145	74.92				
Aug. 2006	5379	7564	71.11				
Nov. 2006	2173	3766	57.7				




Figure 16 : Linear forecast trendline for the dataset of Chak 44/KB Processed by the author

IV. DISCUSSION AND CONCLUSIONS

As the use of space and computer technology developed, humankind has a great advantage to produce this much important research projects with the help of technology in an easier, more accurate way within less time than of other ways. As a result, all these can have a very effective role in helping the country to increase the amount and the quality of agricultural products (Akkartala et al., 2004; Ahmad, 2012; Ahmad, 2012e). With the continuous development of computer network, space technology and remote sensing, demand on automated, real-time and in-orbit

processing become more urgent than ever before for change detection (Mouat et al., 1993; Ahmad, 2012d). To achieve this, the challenges to the technology include the full automation for image registration, image matching, feature extraction, image interpretation, image fusion, data-cleaning, image classification, and data mining and knowledge discovery from GIS database (Liu and Zhou, 2004; Jianya et al., 2008; Ahmad, 2012d). The vegetation indices incorporate a ratio of NIR and red bands, with NDVI being the most frequently used because of its simplicity and robustness. Even so, this index performed relatively poorly for comparison

across our multiple year dataset (Baugh and Groeneveld, 2006).

The NDVI exhibits scaling problems, asymptotic signals over high biomass conditions, and is very sensitive to canopy background variations with NDVI values particularly high with darker canopy backgrounds (Vermote and Vermeulen, 1999; Vermote et al., 2002; Ahmad, 2012b).

RS data can be used to evaluate model performance (Kite and Droogers, 2000; Vazifiedoust et al., 2009), or to initialize, drive, update or re-calibrate models (Droogers and Bastiaanssen, 2002; Vazifiedoust et al., 2009). When the error between model results and data is low, RS-based initialization of models would, in principle, be sufficient to provide spatially distributed information (Vazifiedoust et al., 2009).

Roth (2002) studied the effect of water stress and soil compaction on cotton canopy and showed that canopy reflectance is affected by water stress and canopy reflectance changes were greatest at 555 nm in the green band, 670 nm in the red band and 760 nm in the near infrared band (Kulkarni and Bajwa, 2005). The NDVI can more accurately represent the yield when using higher spatial and temporal resolution satellite data (Li et al., 2007) and the pixel-based modeling revealed a general spatial trend of higher cotton yield (Ruecker et al., 2007).

The methodology presented in this research paper has several desirable properties. Since it treats each pixel individually without setting thresholds or empirical constants, the method is globally applicable (Vermote and Vermeulen, 1999; Vermote et al., 2002; Ahmad, 2012c). The overall accuracy of the results exceeded 85%, the level that the US Geological Survey (Anderson et al., 1976; Sader et al., 2001; Ahmad, 2013a) uses as a threshold to define acceptability (Sader et al., 2001; Ahmad, 2013a). This study also identified several data acquisitions and processing issues that warrant further investigation. Studies are under way to assess the importance of coordinating and timing field data collection and image acquisition dates as a means of improving the strength of the relationships between image and land condition trend analysis (Senseman et al., 1996; Ahmad, 2012e) ground-truth data.

V. ACKNOWLEDGEMENT

The author wishes to thank his respected mentor Dr. Robert Bryant, Sheffield Centre for International Drylands Research, Department of Geography, The University of Sheffield for review and providing valuable comments on draft-version of this paper.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Ackerman, S.A., Strabala, K.I., Menzel, W.P., Frey, R.A., Moeller, C.C., Gumley, L.E., Baum, B., Seemann, S.W. and Hong, Z. (2006). Discriminating clear sky from clouds with MODIS. Algorithm theoretical basis document (MOD35). URL: <http://modis.gsfc.nasa.gov/> (Accessed on December 14, 2012).
2. Ackerman, S.A., Strabala, K.I., Menzel, W.P., Frey, R.A., Moeller, C.C. and Gumley, L.E. (1998). Discriminating clear sky from clouds with MODIS. *Journal of Geophysical Research*, Vol. 103(D24), pp.32,141-32,157.
3. Aguilar, C., Zinnert, J.C., Polo, M.J. and Young, D.R. (2012). NDVI as an indicator for changes in water availability to woody vegetation. *Ecological Indicators*, Vol. 23, pp.290-300.
4. Ahl, D.E., Gower, S.T., Burrows, S.N., Shabanov, N.V., Myneni, R.B. and Knyazikhin, Y. (2006). Monitoring spring canopy phenology of a deciduous broadleaf forest using MODIS. *Remote Sensing of Environment*, Vol. 104, pp.88-95.
5. Ahmad, F. (2012). Phenologically-tuned MODIS NDVI-based time series (2000-2012) for monitoring of vegetation and climatic change in North-Eastern Punjab, Pakistan. *Global Journal of Human Social Science: Geography & Environmental Geo-Sciences*, Vol. 12(13), pp.37-54.
6. Ahmad, F. (2012a). NOAA AVHRR NDVI/MODIS NDVI predicts potential to forest resource management in Çatalca district of Turkey. *Global Journal of Science Frontier Research: Environment & Earth Sciences*, Vol. 12(3), pp.29-46.
7. Ahmad, F. (2012b). NOAA AVHRR satellite data for evaluation of climatic variation and vegetation change in the Punjab Province, Pakistan. *Journal of Food, Agriculture & Environment*, Vol. 10(2), pp.1298-1307.
8. Ahmad, F. (2012c). Landsat ETM+ and MODIS EVI/NDVI data products for climatic variation and agricultural measurements in Cholistan Desert. *Global Journal of Human Social Science: Geography & Environmental Geo-Sciences*, Vol. 12(13), pp.1-11.
9. Ahmad, F. (2012d). A review of remote sensing data change detection: Comparison of Faisalabad and Multan Districts, Punjab Province, Pakistan. *Journal of Geography and Regional Planning*, Vol. 5(9), pp.236-251.

10. Ahmad, F. (2012e). Spectral vegetation indices performance evaluated for Cholistan Desert. *Journal of Geography and Regional Planning*, Vol. 5(6), pp.165-172.
11. Ahmad, F. (2012f). Pixel Purity Index algorithm and n-Dimensional Visualization for ETM+ image analysis: A case of District Vehari. *Global Journal of Human Social Science: Arts and Humanities*, Vol. 12(15), pp.23-32.
12. Ahmad, F. (2013). Detection of change in vegetation cover using multi-spectral and multi-temporal information for District Sargodha. *Global Journal of Human Social Sciences: Geography & Environmental Geo-Sciences*, Vol. 13(1), pp.17-26.
13. Ahmad, F. (2013a). Land degradation pattern using geo-information technology for Kot Addu, Punjab Province, Pakistan. *Global Journal of Human Social Sciences: Geography & Environmental Geo-Sciences*, Vol. 13(1), pp.1-16.
14. Akkartala, A., Türüdüa, O. and Erbekb, F.S. (2004). Analysis of changes in vegetation biomass using multitemporal and multisensor satellite data. In: *Proceedings of XXXV ISPRS Congress*, 12-23 July 2004, Istanbul, Turkey. URL: <http://www.isprs.org/proceedings/XXXV/congress/yf/papers/946.pdf>. (Accessed on September 17, 2011).
15. Al-Awadhi, T., Al-Shukili, A. and Al-Amri, Q. (2011). The use of remote sensing & geographical information systems to identify vegetation: The case of Dhofar Governorate (Oman). URL: <http://www.isprs.org/proceedings/2011/ISRSE-34/211104015Final00239.pdf>. (Assessed on September 10, 2011).
16. Alcaraz-Segura, D., Cabello, J., Paruelo, J.M. and Delibes, M. (2009). Use of descriptors of ecosystem functioning for monitoring a national park network: A remote sensing approach. *Environmental Management*, Vol. 43(1), pp.38-48.
17. Alexandridis, T. and Chemin, Y. (2002). Landsat ETM+, Terra MODIS and NOAA AVHRR: Issues of scale and inter-dependency regarding land parameters. In: *Proceedings of 23rd Asian Conference of Remote Sensing*, 25-29 November 2002, Kathmandu, Nepal, 6 p. URL: <http://www.a-a-r-s.org/acrs/proceeding/ACRS2002/Papers/EOS02-3.htm>. (Accessed on September 07, 2013).
18. Anderson, J.R., Hardy, E.E., Roach, J.T. and Whitmer, R.E. (1976). A land use and land cover classification system for use with remote sensor data. US Geological Survey: Professional Paper 964, Washington, DC.
19. Asrar, G., Fuch, M. and Kanemasu, E.T. (1984). Estimating absorbed photosynthetic radiation and leaf area index from spectral reflectance in wheat. *Agronomy Journal*, Vol. 76, pp.300-306.
20. Asrar, G., Myneni, R.B. and Choudhury, B.J. (1992). Spatial heterogeneity in vegetation canopies and remote sensing of absorbed photosynthetically active radiation: A modeling study. *Remote Sensing of Environment*, Vol. 41, pp. 85-103.
21. Badhwar, G.B. (1984). Automatic corn-soybean classification using Landsat MSS data. II. Early season crop proportion estimation. *Remote Sensing of Environment*, Vol. 14(1-3), pp. 31-37.
22. Baldi, G., Noretto, M.D., Aragón, R., Aversa, F., Paruelo, J.M. and Jobbágy, E.G. (2008). Long-term satellite NDVI datasets: Evaluating their ability to detect ecosystem functional changes in South America. *Sensors*, Vol. 8, pp. 5397-5425.
23. Baret, F. and Guyot, G. (1991). Potentials and limits of vegetation indices for LAI and APAR assessment. *Remote Sensing of Environment*, Vol. 35, pp. 161-173.
24. Barnes, E.M., Moran, M.S., Pinter, Jr. P.J. and Clarke, T.R. (1996). Multispectral remote sensing and site-specific agriculture: Examples of current technology and future possibilities. In: *Proceedings of the 3rd International Conference on Precision Agriculture*, 23-26 June 1996, Minneapolis, Minnesota, USA, pp.843-854.
25. Barrow, C.J. (1991). *Land Degradation: Development and breakdown of terrestrial environments*. Cambridge University Press, Cambridge, p.295.
26. Baugh, W.M. and Groeneveld, D.P. (2006). Broadband vegetation index performance evaluated for a low-cover environment. *International Journal of Remote Sensing*, Vol. 27(21), pp. 4715-4730.
27. Bogue, A. (1993). Leaf area index, intercepted photosynthetically active radiation, and spectral vegetation indices: A sensitivity analysis for regular-clumped canopies. *Remote Sensing of Environment*, Vol. 46, pp.45-59.
28. Beurs, de K.M. and Henebry, G.M. (2004). Land surface phenology, climatic variation, and institutional change: Analyzing agricultural land

cover change in Kazakhstan. *Remote Sensing of Environment*, Vol. 89, pp.497-509.

29. Booth, A., McCullum, J., Mpinga, J. and Mukute, M. (1994). State of the environment in Southern Africa. Southern African Research and Documentation Centre (SARDC), Harare, Zimbabwe. URL: http://rmportal.net/library/content/frame/opinion3.pdf/at_download/file. (Accessed on January 20, 2013).
30. Budde, M.E., Tappan, G., Rowland, J., Lewis, J. and Tieszen, L.L. (2004). Assessing land cover performance in Senegal, West Africa using 1-km integrated NDVI and local variance analysis. *Journal of Arid Environments*, Vol. 59 (3), pp. 481-498.
31. Carrão, H., Gonçalves, P. and Caetano, M. (2008). Contribution of multispectral and multitemporal information from MODIS images to land cover classification. *Remote Sensing of Environment*, Vol. 112(3), pp.986-997.
32. Chen, P.Y., Fedosejevs, G., Tiscareño-López, M. and Arnold, J.G. (2006). Assessment of MODIS-EVI, MODIS-NDVI and VEGETATION-NDVI composite data using agricultural measurements: An example at corn fields in western Mexico. Springer, Netherlands, *Environmental Monitoring and Assessment*, Vol. 119(1-3), pp.69-82.
33. Chen, P.Y., Srinivasan, R., Fedosejevs, G., Báez-González, A.D. and Gong, P. (2002). Assessment of NDVI composite using merged NOAA-14 and NOAA-15 AVHRR data. *Geographic Information Sciences*, Vol. 8(1), pp.31-38.
34. Cheng, Q. (2006). Multisensor comparisons for validation of MODIS vegetation indices. *Pedosphere*, Vol. 16(3), pp.362-370.
35. Cheng, Q. and Wu, X. (2011). Mapping paddy rice yield in Zhejiang Province using MODIS spectral index. *Turkish Journal of Agriculture & Forestry*, Vol. 35(6), pp.579-589.
36. Cheng, Q., Huang, J.F. and Wang, R.C. (2004). Assessment of rice fields by GIS/GPS-supported classification of MODIS data. *Journal of Zhejiang University SCIENCE A: Bioscience & Biotechnology*, Vol. 5(4), pp.412-417.
37. Choudhury, B.J. (1987). Relationship between vegetation indices, radiation absorption, and net photosynthesis evaluated by sensitivity analysis. *Remote Sensing of Environment*, Vol. 22, pp.209-233.
38. Chris, C.F. and Molly, E.B. (2006). Intra-seasonal NDVI change projections in semi-arid Africa. *Remote Sensing of Environment*, Vol. 101, pp.249-256.
39. Clevers, J.G. (1988). The derivation of a simplified reflectance model for the estimation of leaf area index. *Remote Sensing of Environment*, Vol. 25, pp.53-69.
40. Colwell, J.E. (1973). Bidirectional spectral reflectance of grass canopies for determination of above ground standing biomass. Ph.D. Dissertation, University of Michigan, USA.
41. Dabrowska-Zielinska, K., Kogan, F., Ciolkosz, A., Gruszczynska, M. and Kowalik, W. (2002). Modelling of crop growth conditions and crop yield in Poland using AVHRR-based indices. *International Journal of Remote Sensing*, Vol. 23, pp.1109-1123.
42. Dalezios, N.R., Domenikiotis, C., Loukas, A., Tzortzios, S.T. and Kalaitzidis, C. (2001). Cotton yield estimation based on NOAA/AVHRR produced NDVI. *Physics and Chemistry of the Earth, Part B: Hydrology, Oceans and Atmosphere*, Vol. 26(3), pp.247-251.
43. Debien, A., Neerinckx, S., Kimaro, D. and Gulink, H. (2010). Influence of satellite-derived rainfall patterns on plague occurrence in northeast Tanzania. *International Journal of Health Geographics*, Vol. 9, pp.1-10.
44. DeFries, R.S., Hansen, M.C. and Townshend, J.R.G. (1995). Global discrimination of land cover types from metrics derived from AVHRR pathfinder data. *Remote Sensing of Environment*, Vol. 54(3), pp.209-222.
45. Dennison, P.E. and Roberts, D.A. (2003). The effects of vegetation phenology on endmember selection and species mapping in southern California chaparral. *Remote Sensing of Environment*, Vol. 87(2-3), pp.123-135.
46. Dong, J., Kaufmann, R.K., Myneni, R.B., Tucker, C.J., Kauppi, P.E., Liski, J., Buermann, W., Alexeyev, V. and Hughes, M.K. (2003). Remote sensing estimates of boreal and temperate forest woody biomass, carbon pools, sources and sinks. *Remote Sensing of Environment*, Vol. 84, pp. 393-410.
47. Dorman, J.L. and Sellers, P.J. (1989). A Global climatology of albedo, roughness length and stomatal resistance for atmospheric general

- circulation models as represented by the simple biosphere model (SiB). *Journal of Applied Meteorology*, Vol. 28, pp.833-855.
48. Droogers, P. and Bastiaanssen, W.G.M. (2002). Irrigation performance using hydrological and remote sensing modelling. *Journal of Irrigation and Drainage Engineering*, Vol. 128, pp.111-118.
 49. Eastman, J.R. (2003). *Idrisi Kilimanjaro: Guide to GIS and Image Processing*. Clark Labs, Clark University, Worcester, MA. URL: http://www.gis.unbc.ca/help/software/idrisi/kilimanjaro_manual.pdf. (Accessed on January 20, 2013).
 50. Epiphanio, J.C.N. and Huete, A.R. (1995). Dependence of NDVI and SAVI on sun/sensor geometry and its effect on fPAR relationships in Alfalfa. *Remote Sensing of Environment*, Vol. 51, pp. 351-360.
 51. ERDAS Imagine, Inc. (2010). *Professional Tour Guides*. URL: <http://geospatial.intergraph.com/products/ERDASIMAGINE/ERDASIMAGINE/Downloads.aspx>. (Accessed on February 18, 2013).
 52. Fensholt, R. (2004). Earth observation of vegetation status in the Sahelian and Sudanian West Africa: Comparison of Terra MODIS and NOAA AVHRR satellite data. *International Journal of Remote Sensing*, Vol. 10, pp.1641-1659.
 53. Fontana, F.M.A. (2009). From single pixel to continental scale: using AVHRR and MODIS to study land surface parameters in mountain regions. Ph.D. dissertation, Institute of Geography, University of Bern, Switzerland. URL: <http://www.climatestudies.unibe.ch/students/theses/phd/31.pdf>. (Accessed on October 15, 2012).
 54. Galio, K.P., Daughtry, C.S.T. and Bauer, M.E. (1985). Spectral estimation of absorbed photosynthetically active radiation in corn canopies. *Agronomy Journal*, Vol. 78, pp.752-756.
 55. Gao, B-C. (1996). NDWI - A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment*, Vol. 58(3), pp.257-266.
 56. Gao, X., Huete, A.R., Ni, W. and Miura, T. (2000). Optical-biophysical relationships of vegetation spectra without background contamination. *Remote Sensing of Environment*, Vol. 74(3), pp. 609-620.
 57. Gillies, R.R., Carlson, T.N., Cui, J., Kustas, W.O. and Humes, K.S. (1997). A verification of the 'triangle' method for obtaining surface soil water content and energy fluxes from remote measurements of the Normalized Difference Vegetation Index (NDVI) and surface radiant temperature. *International Journal of Remote Sensing*, Vol. 18, pp.3145-3166.
 58. Gitelson, A.A. and Merzlyak, M.N. (1996). Signature analysis of leaf reflectance spectra: Algorithm development for remote sensing of chlorophyll. *Journal of Plant Physiology*, Vol. 148, pp.494-500.
 59. Glenn, E.P., Huete, A.R., Nagler, P.L. and Nelson, S.G. (2008). Relationship between remotely-sensed vegetation indices, canopy attributes and plant physiological processes: What vegetation indices can and cannot tell us about the landscape. *Sensors*, Vol. 8, pp.2136-2160.
 60. Groeneveld, D.P. and Baugh, W.M. (2007). Correcting satellite data to detect vegetation signal for eco-hydrologic analyses. *Journal of Hydrology*, Vol. 344(1-2), pp.135-145.
 61. GSFC/NASA (2003). MODIS technical specifications. URL: <http://modis.gsfc.nasa.gov/about/specs.html>. (Accessed on September 05, 2011).
 62. Hall, F.G. and Badhwar, G.D. (1987). Signature-extendable technology: Global space-based crop recognition. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. GE-25(1), pp.93-103.
 63. Hamel, S., Garel, M., Festa-Bianchet, M., Gaillard, J-M. and Côté, S.D. (2009). Spring Normalized Difference Vegetation Index (NDVI) predicts annual variation in timing of peak faecal crude protein in mountain ungulates. *Journal of Applied Ecology*, Vol. 46(3), pp. 582-589.
 64. Hobbs, R.J. (1989). Remote sensing of spatial and temporal dynamics of vegetation. In: Hobbs, R.J. and Mooney, H.A. (Eds.), *Remote Sensing of Biosphere Functioning*, Springer-Verlag, New York, pp.203-219.
 65. Holben, B. (1986). Characteristics of maximum-value composite images from temporal AVHRR data. *International Journal of Remote Sensing*, Vol. 7, pp.1417-1434.
 66. Hongo, C. and Niwa, K. (2012). Yield prediction of sugar beet through combined use of satellite data and meteorological data. *Journal of Agricultural Science*, Vol. 4(4), p.251-261.
 67. Höpfner, C. and Scherer, D. (2011). Analysis of vegetation and land cover dynamics in north-western Morocco during the last decade using MODIS NDVI time series data. *Biogeosciences*, Vol. 8, pp.3359-3373.

68. Huang, Q., Li, M., Chen, C., Mao, K., Chen, Z., Li, F. and Chen, D. (2010). Assessment of land degradation in Guizhou province, Southwest China using AVHRR/NDVI and MODIS/NDVI data. 18th International Conference on Geoinformatics, 18-20 June 2010, Beijing, pp.1-5. doi: 10.1109/GEOINFORMATICS.2010.5568119
69. Huete, A.R. (2005). Global variability of terrestrial surface properties derived from MODIS visible to thermal-infrared measurements. In: Proceedings of IGARSS 05 Geoscience and Remote Sensing Symposium, 25-29 July 2005. URL: <http://ieeexplore.ieee.org/iel5/10226/32601/01526782.pdf>. (Accessed on September 05, 2011).
70. Huete, A.R., Didan, K., Miura, T., Rodriguez, E.P., Gao, X. and Ferreira, L.G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment*, Vol. 83(1-2), pp.195-213.
71. Huete, A.R., Didan, K., Shimabukuro, Y.E., Ratana, P., Saleska, S.R., Hutya, L.R., Yang, W., Nemani, R.R. and Myneni, R. (2006). Amazon rainforests green-up with sunlight in dry season. *Geophysical Research Letters*, 33(L06405), 4 p.
72. Huete, A.R., Justice, C.O. and Leeuwen, van W. (1999). MODIS vegetation index (MOD 13) algorithm theoretical basis document version 3. URL: http://modis.gsfc.nasa.gov/data/atbd/atbd_mod13.pdf. (Accessed on September 17, 2011).
73. Huete, A.R., Liu, H.Q., Batchily, K. and Leeuwen, van W. (1997). A comparison of vegetation indices over a global set of TM images for EOS-MODIS. *Remote Sensing of Environment*, Vol. 59, pp. 440-451.
74. Hutchinson, C.F. (1991). Uses of satellite data for famine early warning in sub-Saharan Africa. *International Journal of Remote Sensing*, Vol. 12, pp.1405-1421.
75. Jakubauskas, M.E., David, R. and Kastens, J.H. (2002). Crop identification using harmonic analysis of time-series AVHRR NDVI data. *Computers and Electronics in Agriculture*, Vol. 37(1-3), pp.127-139.
76. Jensen, J.R. (2005). *Introductory digital image processing: A remote sensing perspective*. 3rd Edition, Englewood Cliffs, New Jersey, Prentice-Hall, pp.34-67.
77. Jiang, Z., Huete, A.R., Didan, K. and Miura, T. (2008). Development of a two-band Enhanced Vegetation Index without a blue band. *Remote Sensing of Environment*, Vol. 112(10), pp.3833-3845.
78. Jianya, G., Haigang, S., Guorui, Ma. and Qiming, Z. (2008). A review of multi-temporal remote sensing data change detection algorithms. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, Vol. XXXVII, Part B7, pp.757-762. URL: http://www.isprs.org/proceedings/XXXVII/congress/7_pdf/5_WG-VII-5/05.pdf. (Accessed on November 04, 2011).
79. Jin, S. and Sader, S.A. (2005). MODIS time-series imagery for forest disturbance detection and quantification of patch size effects. *Remote Sensing of Environment*, Vol. 99, pp. 462-470.
80. Johnson, J.F., Voorhees, W.B., Nelson, W.W. and Randall, G.W. (1990). Soybean growth and yield as affected by surface and subsoil compaction. *Agronomy Journal*, Vol. 82(5), pp. 973-979.
81. Justice, C.O., Townshend, J.R.G., Holben, B.N. and Tucker, C.J. (1985). Analysis of the phenology of global vegetation using meteorological satellite data. *International Journal of Remote Sensing*, Vol. 6, pp. 1271-1318.
82. Justice, C.O., Vermote, E., Townshend, J.R.G., DeFries, R.S., Roy, D.P., Hall, D.K., Salomonson, V.V., Privette, J., Riggs, G., Strahler, A., Lucht, W., Myneni, R., Knjazihhin, Y., Running, S., Nemani, R., Wan, Z., Huete, A.R., Leeuwen, van W., Wolfe, R.E., Giglio, L., Muller, J.-P., Lewis, P. and Barnsley, M. (1998). The Moderate Resolution Imaging Spectroradiometer (MODIS): Land remote sensing for global change research. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 36(4), pp. 1228-1249.
83. Karaburun, A. (2010). Estimation of C factor for soil erosion modeling using NDVI in Büyükçekmece watershed. *Ozean Journal of Applied Sciences*, Vol. 3(1), pp. 77-85.
84. Khlopenkov, K.V. and Trishchenko, A.P. (2008). Implementation and evaluation of concurrent gradient search method for reprojection of MODIS level 1B imagery. *IEEE Transaction on Geoscience and Remote Sensing*, Vol. 46(7), pp. 2016-2027.
85. Kite, G.W. and Droogers, P. (2000). Comparing evapotranspiration estimates from satellites, hydrological models and field data. *Journal of Hydrology*, Vol. 229, pp.3-18.

86. Kogan, F.N. (1987). Vegetation index for areal analysis of crop conditions. In: Proceedings of 18th Conference Agricultural and Forest Meteorology and 8th Conference Biometeorology and Aerobiology, 15-18 September 1987, W. Lafayette, Indiana, AMS Publisher, pp.103-106.
87. Korontzi, S., Roy, D.P., Justice, C.O. and Ward, D.E. (2004). Modeling and sensitivity analysis of fire emissions in southern Africa during SAFARI 2000. *Remote Sensing of Environment*, Vol. 92, pp.376-396.
88. Kulkarni, S.S. and Bajwa, S.G. (2005). Spectral response of cotton canopy to soil compaction. Paper Number: 051066, The Society for engineering in agricultural, food, and biological systems, USA, pp.1-19.
89. Lambin, E.F. (1996). Change detection at multiple temporal scales - seasonal and annual variations in landscape variables. *Photogrammetric Engineering & Remote Sensing*, Vol. 62, pp.931-938.
90. Li, A., Liang, S., Wang, A. and Qin, J. (2007). Estimating crop yield from multi-temporal satellite data using multivariate regression and neural network techniques. *Photogrammetric Engineering & Remote Sensing*, Vol. 73(10), pp.1149-1157.
91. Lillesand, T.M. and Kiefer, R.W. (2000). *Remote sensing and image interpretation*. 4th Edition, John Wiley & Sons, Inc., New York, pp.124-156.
92. Lillesand, T.M., Kiefer, R.W. and Chipman, J.W. (2004). *Remote sensing and image interpretation*. 5th Edition, John Wiley & Sons, Inc., New York, pp.178-198.
93. Liu, H. and Zhou, Q. (2004). Accuracy analysis of remote sensing change detection by rule-based rationality evaluation with post-classification comparison. *International Journal of Remote Sensing*, Vol. 25(5), pp.1037-1050.
94. Liu, H.Q. and Huete, A.R. (1995). A feedback based modification of the NDVI to minimize canopy background and atmospheric noise. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 33(2), pp.457-465.
95. Liu, W.T. and Kogan, F. (2002). Monitoring Brazilian soybean production using NOAA/AVHRR based vegetation condition indices. *International Journal of Remote Sensing*, Vol. 23(6), pp.1161-1179.
96. Lloyd, D. (1990). A phenological classification of terrestrial vegetation cover using shortwave vegetation index imagery. *International Journal of Remote Sensing*, Vol. 11(12), pp.2269-2279.
97. Lobell, D.B., Asner, G.P., Ortiz-Monasterio, J.I. and Benning, T.L. (2003). Remote sensing of regional crop production in the Yaqui Valley, Mexico: estimates and uncertainties. *Agriculture, Ecosystems and Environment*, Vol. 94, pp.205-220.
98. Logan, T.L. (1999). EOS/AM-1 Digital Elevation Model (DEM) Data Sets: DEM and DEM auxiliary datasets in support of the EOS/Terra platform. JPL D-013508. Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA.
99. Los, S.O. (1998). Estimation of the ratio of sensor degradation between NOAA-AVHRR channels 1 and 2 from monthly NDVI composites. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 36, pp.206-213.
100. Los, S.O., Collatz, G.J., Bounoua, L., Sellers, P.J. and Tucker, C.J. (2001). Global interannual variations in sea surface temperature and land surface vegetation, air temperature, and precipitation. *Journal of Climate*, Vol. 14, pp. 1535-1549.
101. Los, S.O., Collatz, G.J., Sellers, P.J., Malmström, C.M., Pollack, N.H., DeFries, R.S., Bounoua, L., Parris, M.T., Tucker, C.J. and Dazlich, D.A. (2000). A Global 9-yr biophysical land surface dataset from NOAA- AVHRR data. *Journal of Hydrometeorology*, Vol. 1, pp. 183-199.
102. Los, S.O., Justice, C.O. and Tucker, C.J. (1994). A global 1° by 1° NDVI data set for climate studies derived from the GIMMS continental NDVI data. *International Journal of Remote Sensing*, Vol. 15(17), pp. 3493-3518.
103. Los, S.O., North, P.R.J., Grey, W.M.F. and Barnsley, M.J. (2005). A method to convert AVHRR normalized difference vegetation index time series to a standard viewing and illumination geometry. *Remote Sensing of Environment*, Vol. 99(4), pp. 400-411.
104. Lowery, B. and Schuler, R.T. (1991). Temporal effects of subsoil compaction on soil strength and plant growth. *Soil Science Society of America Journal*, Vol. 55(1), pp.216-223.
105. Lu, D., Mausel, P., Brondizios, E. and Moran, E. (2004). Change detection techniques. *International Journal of Remote Sensing*, Vol. 25(12), pp. 2365-2407.

106. Lyon, J.G., Yuan, D., Lunetta, R.S. and Elvidge, C.D. (1998). A change detection experiment using vegetation indices. *Photogrammetric Engineering & Remote Sensing*, Vol. 64(2), pp. 143-150.
107. Macdonald, R.B. and Hall, F.G. (1980). Global crop forecasting. *Science*, Vol. 208(4445), pp.670-679.
108. Macleod, R.D. and Congalton, R.G. (1998). A quantitative comparison of change detection algorithms for monitoring eelgrass from remotely sensed data. *Photogrammetric Engineering and Remote Sensing*, Vol. 64(3), pp.207-216.
109. Mahmoodzadeh, H. (2007). Digital change detection using remotely sensed data for monitoring green space destruction in Tabriz. *International Journal of Environmental Research*, Vol. 1(1), pp. 35-41.
110. Mambo, J. and Archer, E. (2007). An assessment of land degradation in the Save catchment of Zimbabwe. *Area*, Vol. 39(3), pp. 380-391.
111. Mas, J-F. (1999). Monitoring land-cover changes: a comparison of change detection techniques. *International Journal of Remote Sensing*, Vol. 20(1), pp. 139-152.
112. Meneses-Tovar, C.L. (2011). NDVI as indicator of degradation. *Unasylva*, Vol. 62(238), pp. 39-46.
113. Milich, L. and Weiss, E. (2000). GAC NDVI interannual coefficient of variation (CoV) images: Ground truth sampling of the Sahel along north-south transects. *International Journal of Remote Sensing*, Vol. 21, pp. 235-260.
114. Mirik, M., Michels, Jr. G.J., Kassymzhanova, M.S., Elliott, N.C. and Catana, V. (2006). Spectral sensing of aphid (Hemiptera: Aphididae) density using field spectrometry and radiometry. *Turkish Journal of Agriculture and Forestry*, Vol. 30, pp. 421-428.
115. Mirik, M., Norland, J.E., Biondini, M.E., Crabtree, R.L. and Michels, Jr. G.J. (2007). Relationships between remotely sensed data and biomass components in a big sagebrush (*Artemisia tridentata*) dominated area in Yellowstone national park. *Turkish Journal of Agriculture and Forestry*, Vol. 31, pp. 135-145.
116. MODIS (1999). MODIS Vegetation Index (MOD 13): Algorithm theoretical basis document pp.26-29 (version 3). URL: http://modis.gsfc.nasa.gov/data/atbd/atbd_mod13.pdf. (Accessed on December 04, 2008).
117. Moran, M.S., Inoue, Y. and Barnes, E.M. (1997). Opportunities and limitations for image based remote sensing in precision crop management. *Remote Sensing of Environment*, Vol. 61, pp. 319-346.
118. Moran, M.S., Jackson, R.D., Slater, P.N. and Teillet, P.M. (1992). Evaluation of simplified procedures for retrieval of land surface reflectance factors from satellite sensor output. *Remote Sensing of Environment*, Vol. 41, pp. 169-184.
119. Mouat, D.A., Mahin, G.C. and Lancaster, J. (1993). Remote sensing techniques in the analysis of change detection. *Geocarto International*, Vol. 2, pp. 39-50.
120. Moulin, S., Kergoat, L., Viovy, N. and Dedieu, G. (1997). Global scale assessment of vegetation phenology using NOAA/AVHRR satellite measurements. *Journal of Climate*, Vol. 10, pp. 1154-1170.
121. Myneni, R.B. and Asrar, G. (1994). Atmospheric effects and spectral vegetation indices. *Remote Sensing of Environment*, Vol. 47, pp. 390-402.
122. Myneni, R.B., Dong, J., Tucker, C.J., Kaufmann, R.K., Kauppi, P.E., Liski, J., Zhou, L., Alexeyev, V. and Hughes, M.K. (2001). A large carbon sink in the woody biomass of Northern forests. In: *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 98, pp. 14784-14789.
123. Myneni, R.B., Hall, F.G., Sellers, P.J. and Marshak, A.L. (1995). The interpretation of spectral vegetation indices. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 33(2), pp. 481-486.
124. Myneni, R.B., Keeling, C.D., Tucker, C.J., Asrar, G. and Nemani, R.R. (1997). Increased plant growth in the northern high latitudes from 1981-1991. *Nature*, Vol. 386, pp. 698-702.
125. Nasr, A.H. and Helmy, A.K. (2009). Integration of MIRSAT-1 and SPOT-2 data for quantitative change detection applications. *ICGST-GVIP Journal*, Vol. 9(5), pp.53-59. URL: http://www.icgst.com/gvip/volume9/Issue5/GVIP_V9_I5.pdf. (Accessed on November 04, 2011).
126. Nicandrou, A. (2010). Hydrological assessment and modelling of the river Fani catchment, Albania. Faculty of Advanced Technology, University of Glamorgan. URL: <http://dspace1.isd.glam.ac.uk/dspace/handle/10265/461> (Accessed on November 11, 2012).
127. Nicholson, S.E. and Farrar, T. (1994). The influence of soil type on the relationships between NDVI,

- rainfall and soil moisture in semi-arid Botswana: Part I. NDVI response to rainfall. *Remote Sensing of Environment*, Vol. 50, pp.107-120.
128. Nicholson, S.E., Tucker, C.J. and Ba, M.B. (1998). Desertification, drought, and surface vegetation: An example from the West African Sahel. *Bulletin of the American Meteorological Society*, Vol. 79, pp. 1-15.
 129. Ouyang, W., Hao, F.H., Skidmore, A.K., Groen, T.A., Toxopeus, A.G. and Wang, T. (2012). Integration of multi-sensor data to assess grassland dynamics in a Yellow River sub-watershed. *Ecological Indicators*, Vol. 18, pp. 163-170.
 130. Paruelo, J.M., Epstein, H.E., Lauenroth, W.K. and Burke, I.C. (1997). ANPP estimates from NDVI for the Central Grassland Region of the United States. *Ecology*, Vol. 78(3), pp. 953-958.
 131. Pettorelli, N., Vik, J.O., Mysterud, A., Gaillard, J.M., Tucker, C.J. and Stenseth, N.C. (2005). Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution*, Vol. 20(9), pp. 503-510.
 132. Piao, S.L., Mohammat, A., Fang, J.Y., Cai, Q. and Feng, J. (2006). NDVI-based increase in growth of temperate grasslands and its responses to climate changes in China. *Global Environmental Change*, Vol. 16(4), pp. 340-348.
 133. Piao, S.L., Fang, J.Y., Zhou, L.M., Guo, Q.H., Henderson, M., Ji, W., Li, Y. and Tao, S. (2003). Interannual variations of monthly and seasonal NDVI in China from 1982 to 1999. *Journal of Geophysical Research*, Vol. 108(D14), pp. 4401-4413.
 134. Potter, C.S., Randerson, J.T., Field, C.B., Matson, P.A., Vitousek, P.M., Mooney, H.A. and Klooster, S.A. (1993). Terrestrial ecosystem production, a process model based on global satellite and surface data. *Global Biogeochemical Cycles*, Vol. 7, pp. 811-841.
 135. Prasad, A.K., Sudipta, S., Singh, R.P. and Kafatos, M. (2007). Inter-annual variability of vegetation cover and rainfall over India. *Advances in Space Research*, Vol. 39(1), pp.79-87.
 136. Price, J.C. and Bausch, W.C. (1995). Leaf area index estimation from visible and NIR-infrared reflectance data. *Remote Sensing of Environment*, Vol. 52, pp. 55-65.
 137. Price, K.P., Egbert, S.L., Nellis, M.D., Lee, R-Y. and Boyce, R. (1997). Mapping land cover in a high plains agro-ecosystem using a multirate Landsat thematic mapper modeling approach. *Transactions of the Kansas Academy of Science*, Vol. 100(1-2), pp. 21-33.
 138. Prince, S.D. (1991). Satellite remote sensing of primary production: Comparison of results for Sahelian grasslands 1981-1988. *International Journal of Remote Sensing*, Vol. 12, pp.1301-1311.
 139. Prince, S.D. and Tucker, C.J. (1986). Satellite remote sensing of rangelands in Botswana: II. NOAA AVHRR and herbaceous vegetation. *International Journal of Remote Sensing*, Vol. 7, pp.1555-1570.
 140. Qi, J., Chehbouni, A., Huete, A.R., Kerr, Y.H. and Sorooshian, S. (1994). A modified soil adjusted vegetation index. *Remote Sensing of Environment*, Vol. 48, pp. 119-126.
 141. Rahman, Md.R., Islam, A.H.M.H. and Rahman, Md.A. (2004). NDVI derived sugarcane area identification and crop condition assessment. *Planplus*, Vol. 1(2), pp. 1-12.
 142. Reed, B.C., Brown, J.F., VanderZee, D., Loveland, T.R., Merchant, J.W. and Ohlen, D.O. (1994). Measuring phenological variability from satellite imagery. *Journal of Vegetation Science*, Vol. 5(5), pp.703-714.
 143. Research System, Inc. (2004). Exploring ENVI, Boulder, CO. URL: <http://www.itvis.com> (Accessed on November 28, 2011).
 144. Richard, J., Radke, S.A., Omar, A.K. and Badrinath, R. (2005). Image change detection algorithms: A systematic survey. *IEEE Transactions on Image Processing*, Vol. 14(3), pp. 294-307.
 145. Rosenthal, W.D. and Gerik, T.J. (1991). Radiation use efficiency among cotton cultivars. *Agronomy Journal*, Vol. 83(4), pp. 655-658.
 146. Roth, G.W. (2002). The effect of water stress and soil compaction on cotton canopy reflectance and temperature. In: *Proceedings of the 11th Cotton Conference*, Brisbane, Australia, pp.623-630.
 147. Rouse, J.W., Haas, R.H., Schell, J.A. and Deering, D.W. (1973). Monitoring vegetation systems in the Great Plains with ERTS. 3rd ERTS Symposium, NASA SP-351 I, pp. 309-317. URL: http://geol.hu/data/online_help/Vegetation_Indices.html. (Accessed on January 10, 2012).
 148. Roy, D.P., Lewis, P.E. and Justice, C.O. (2002). Burned area mapping using multi-temporal moderate spatial resolution data – a bi-directional reflectance model-based expectation approach.

- Remote Sensing of Environment, Vol. 83(1-2), pp. 263-286.
149. Ruecker, G.R., Shi, Z., Mueller, M., Conrad, C., Ibragimov, N., Lamers, J.P.A., Martius, C., Strunz, G. and Dech, S.W. (2007). Cotton yield estimation in Uzbekistan integrating MODIS, Landsat ETM+ and field data. pp.123-128. URL: http://www.isprs.org/proceedings/xxxvi/8-w48/123_XXXVI-8-W48.pdf. (Accessed on September 18, 2013).
 150. Running, S.W. and Nemani, R.R. (1991). Regional hydrologic and carbon balance responses of forests resulting from potential climate change. *Climatic Change*, Vol. 19, pp. 349-368.
 151. Running, S.W., Justice, C.O., Salomonson, V.V., Hall, D., Barker, J., Kaufman, Y.J., Strahler, A.H., Huete, A.R., Muller, J.-P., Vanderbilt, V., Wan, Z.M., Teillet, P. and Carneggie, D. (1994). Terrestrial remote sensing science and algorithms planned for EOS/MODIS. *International Journal of Remote Sensing*, Vol. 15, pp. 3587-3620.
 152. Sader, S.A., Hayes, D.J., Hepinstall, J.A., Coan, M. and Soza, C. (2001). Forest change monitoring of a remote biosphere reserve. *International Journal of Remote Sensing*, Vol. 22(10), pp.1937-1950.
 153. Salomonson, V.V. and Toll, D.L. (1991). Execution phase (C/D) spectral band characteristics of the EOS moderate resolution imaging spectrometer-Nadir (MODIS-N) facility instrument. *Advances in Space Research*, Vol. 11, pp.231-236.
 154. Schroeder, T.A., Canty, M.J. and Yang, Z. (2006). Radiometric correction of multi-temporal Landsat data for characterization of early successional forest patterns in western Oregon. *Remote Sensing of Environment*, Vol. 103, pp. 16-26.
 155. Sellers, P.J. (1985). Canopy reflectance, photosynthesis and transpiration. *International Journal of Remote Sensing*, Vol. 6, pp.1335-1372.
 156. Sellers, P.J., Los, S.O., Tucker, C.J., Justice, C.O., Dazlich, D.A., Collatz, G.J. and Randall, D.A. (1996). A revised land surface parameterization (SiB-2) for atmospheric GCMs. Part 2: The generation of global fields of terrestrial biophysical parameters from satellite data. *Journal of Climate*, Vol. 9(4), pp. 706-737.
 157. Sellers, P.J., Tucker, C.J., Collatz, G.J., Los, S.O., Justice, C.O., Dazlich, D.A. and Randall, D.A. (1994). A global 1o by 1o NDVI data set for climate studies, Part 2: The adjustment of the NDVI and generation of global fields of terrestrial biophysical parameters. *International Journal of Remote Sensing*, Vol. 15, pp.3519-3545.
 158. Senseman, G.M., Bagley, C.F. and Tweddale, S.A. (1996). Correlation of rangeland cover measures to satellite-imagery-derived vegetation indices. *Geocarto International*, Vol. 11(3), pp.29-38.
 159. Sesnie, S.E., Gessler, P.E., Finegan, B. and Thessler, S. (2008). Integrating Landsat TM and SRTM-DEM derived variables with decision trees for habitat classification and change detection in complex neotropical environments. *Remote Sensing of Environment*, Vol. 112, pp.2145-2159.
 160. Shi, Z., Ruecker, G.R., Mueller, M., Conrad, C., Ibragimov, N., Lamers, J.P.A., Martius, C., Strunz, G., Dech, S. and Vlek, P.L.G. (2007). Modeling of cotton yields in the Amu Darya river floodplains of Uzbekistan integrating multitemporal remote sensing and minimum field data. *Agronomy Journal*, Vol. 99(5), pp.1317-1326.
 161. Shinnars, K.J. and Binversie, B.N. (2007). Fractional yield and moisture of corn stover biomass produced in the Northern US Corn Belt. *Biomass and Bioenergy*, Vol. 31, pp.576-584.
 162. Shippert, P. (2004). Why use hyperspectral imagery? *Photogrammetric Engineering & Remote Sensing*, pp.377-380. URL: <http://www.iro.umontreal.ca/~mignotte/IFT6150/ComplementCours/HyperspectralImagery.pdf>. (Accessed on November 20, 2011).
 163. Singh, A. (1989). Digital change detection techniques using remotely sensed data. *International Journal of Remote Sensing*, Vol. 10(6), pp. 989-1003.
 164. Singh, D. (2012). Evaluation of long-term NDVI time series derived from Landsat data through blending with MODIS data. *Atmósfera*, Vol. 25(1), pp.43-63.
 165. Singh, R.P., Roy, S. and Kogan, F. (2003). Vegetation and temperature condition indices from NOAA AVHRR data for drought monitoring over India. *International Journal of Remote Sensing*, Vol. 24(22), pp. 4393-4402.
 166. Smith, J.H. and Ramey, D.B. (1982). A crop area estimator based on changes in the temporal profile of a vegetative index. In: *Proceedings of the Survey Research Methods Section, American Statistical Association*, Cincinnati, Ohio, USA, pp.495-498. URL: http://www.amstat.org/sections/srms/Proceedings/papers/1982_095.pdf. (Accessed on September 21, 2013).

- Vegetation and temperature condition indices from NOAA AVHRR data for drought monitoring over India. *International Journal of Remote Sensing*, Vol. 24(22), pp. 4393-4402.
167. Sönmez, N.K. and Sari, M. (2006). Use of remote sensing and geographic information system technologies for developing greenhouse databases. *Turkish Journal of Agriculture and Forestry*, Vol. 30, pp. 413-420.
 168. Stow, D. (1995). Monitoring ecosystem response to global change: Multitemporal remote sensing analyses. In: Moreno, J. and Oechel, W. (Eds.), *Anticipated Effects of a Changing Global Environment in Mediterranean Type Ecosystems*, Springer-Verlag, New York, pp.254-286.
 169. Symeonakis, E., Caccetta, P.A., Wallace, J.F. and Koukoulas, S. (2006). Multi-temporal land use/cover change detection in the Spanish Mediterranean coast. In: *Proceedings of ISPRS Mid-Term Symposium, Remote Sensing: From pixels to processes*, ITC, The Netherlands, pp.58-63. URL: <http://www.isprs.org/proceedings/XXXVI/part7/PDF/058.pdf>. (Accessed on November 04, 2011).
 170. Tarpley, J.D., Schnieder, S.R. and Money, R.L. (1984). Global vegetation indices from NOAA-7 meteorological satellite. *Journal of Climate and Applied Meteorology*, Vol. 23, pp.4491-4503.
 171. Taylor, H.M. and Gardner, H.R. (1963). Penetration of cotton seedlings tap roots as influenced by bulk density, moisture content and strength of soil. *Soil Science*, Vol. 96, pp.153-156.
 172. Thenkabail, P.S., Smith, R.B. and Eddy De Pauw, E. De. (2000). Hyperspectral vegetation indices and their relationships with agricultural crop characteristics. *Remote Sensing of Environment*, Vol. 71, pp.158-182.
 173. Thiam, A. (2003). The causes and spatial pattern of land degradation risk in southern Mauritania using multitemporal AVHRR-NDVI imagery and field data. *Land Degradation & Development*, Vol. 14, pp. 133-142.
 174. Thomas, D.S.G. and Leason, H.C. (2005). Dunefield activity response to climate variability in the southwest Kalahari. *Geomorphology*, Vol. 64(1-2), pp.117-132.
 175. Townshend, J.R.G. (1992). Improved global data for land applications: A proposal for a new high resolution data set. Report No. 20. International Geosphere-Biosphere Program, Stockholm, Sweden. URL: <http://library.wur.nl/WebQuery/clc/916308>. (Accessed on June 04, 2012).
 176. Townshend, J.R.G. and Justice, C.O. (1988). Selecting the spatial-resolution of satellite sensors required for global monitoring of land transformations. *International Journal of Remote Sensing*, Vol. 9(2), pp.187-236.
 177. Townshend, J.R.G., Justice, C.O., Li, W., Gurney, C. and McManus, J. (1991). Global land cover classification by remote sensing: present capabilities and future possibilities. *Remote Sensing of Environment*, Vol. 35(2-3), pp.243-255.
 178. Tucker, C.J. (1979). Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment*, Vol. 8, pp.127-150.
 179. Tucker, C.J., Dregne, H.E. and Newcomb, W.W. (1991). Expansion and contraction of the Sahara Desert from 1980 to 1990. *Science*, Vol. 253, pp.299-301.
 180. Tucker, C.J., Gatlin, J., Schnieder, S.R. and Kuchinos, M.A. (1982). Monitoring large scale vegetation dynamics in the Nile delta and river valley from NOAA AVHRR data. In: *Proceedings of the Conference on Remote Sensing of Arid and Semi-Arid Lands*, Cairo, Egypt, Ann Arbor, MI. pp.973-977.
 181. Ulbricht, K.A. and Heckendorf, W.D. (1998). Satellite images for recognition of landscape and land use changes. *ISPRS Journal of Photogrammetry & Remote Sensing*, Vol. 53, pp.235-243.
 182. Unger, P.W. and Kasper, T.C. (1994). Soil compaction and root growth: A review. *Agronomy Journal*, Vol. 86(5), pp.759-766.
 183. USGS (2008). Earth Resources Observation and Science Center. URL: <http://glovis.usgs.gov/> (Accessed on September 09, 2013).
 184. USGS (2010). What is NDVI? United States Geological Survey: Science for Changing World. URL: <http://ivm.cr.usgs.gov/> (Accessed on September 10, 2011).
 185. Vazifedoust, M., Van Dam, J.C., Bastiaanssen, W.G.M. and Feddes, R.A. (2009). Assimilation of satellite data into agrohydrological models to improve crop yield forecasts. *International Journal of Remote Sensing*, Vol. 30(10), pp.2523-2545.
 186. Verbesselt, J., Hyndman, R., Zeileis, A. and Culvenor, D. (2010). Phenological change detection while accounting for abrupt and gradual

- trends in satellite image time series. *Remote Sensing of Environment*, Vol. 114(12), pp.2970-2980.
187. Vermote, E.F. and Vermeulen, A. (1999). Atmospheric correction algorithm: Spectral reflectances (MOD09) algorithm technical background document. URL: http://modis.gsfc.nasa.gov/data/atbd/atbd_mod08.pdf. (Accessed on September 07, 2011).
 188. Vermote, E.F., El-Saleous, N.Z. and Justice, C.O. (2002). Atmospheric correction of MODIS data in the visible to middle infrared: First results. *Remote Sensing of Environment*, Vol. 83, pp.97-111.
 189. Wan, Z. (2003). MODIS Land-Surface Temperature products users' guide. URL: <http://www.ices.ucsb.edu/modis/LstUsrGuide/usrguide.html>. (Accessed on September 07, 2011).
 190. Wang, J., Rich, P.M. and Price, K.P. (2003). Temporal responses of NDVI to precipitation and temperature in the central Great Plains, USA. *International Journal of Remote Sensing*, Vol. 24, pp.2345-2364.
 191. Wardlow, B.D., Egbert, S.L. and Kastens, J.H. (2007). Analysis of time-series MODIS 250 m vegetation index data for crop classification in the US Central Great Plains. *Remote Sensing of Environment*, Vol. 108(3), pp.290-310.
 192. Wessels, K.J., Prince, S.D., Frost, P.E. and Zyl, van D. (2004). Assessing the effects of human-induced land degradation in the former homelands of northern South Africa with a 1 km AVHRR NDVI time-series. *Remote Sensing of Environment*, Vol. 91, pp.47-67.
 193. Wiegant, C.L., Richardson, A.J. and Kanemasu, K. (1979). Leaf area index estimates for wheat from Landsat and their implications for evapotranspiration and crop modeling. *Agronomy Journal*, Vol. 71, pp.336-342.
 194. Wolfe, R.E., Storey, J., Masuoka, E. and Fleig, A. (1995). MODIS Level 1A Earth location algorithm theoretical basis document. URL: <http://modis.gsfc.nasa.gov/> (Accessed on December 14, 2012).
 195. Wolfe, R.E., Nishihama, M., Fleig, A.J., Kuyper, J.A., Roy, D.P., Storey, J.C. and Patt, F.S. (2002). Achieving sub-pixel geolocation accuracy in support of MODIS land science. *Remote Sensing of Environment*, Vol. 83(1-2), pp.31-49.
 196. Xie, Y. (2008). Remote sensing imagery in vegetation mapping: A review. *Journal of Plant Ecology*, Vol. 1(1), pp.9-23.
 197. Yang, Y., Yang, L. and Merchant, J.W. (1997). An assessment of AVHRR/NDVI-ecoclimatological relations in Nebraska, USA. *International Journal of Remote Sensing*, Vol. 18(10), pp.2161-2180.
 198. Young, A. (1998). Assessment of land degradation by remote sensing in land resources: Now and for the Future. Cambridge University Press, Cambridge. URL: <http://www.uea.ac.uk/env/landresources/news-degradation-rs.html>. (Accessed on January 20, 2013).
 199. Zaeen, A.A. (2012). Remote sensing technique to monitoring the risk of soil degradation using NDVI. *International Journal of Geographical Information System Applications and Remote Sensing*, Vol. 3(1), pp.1-8.
 200. Zhan, X., DeFries, R.S., Townshend, J.R.G., DiMiceli, C., Hansen, M.C., Huang, C. and Sohlberg, R.A. (2000). The 250m global land cover change product from the Moderate Resolution Imaging Spectroradiometer of NASA's Earth Observing System. *International Journal of Remote Sensing*, Vol. 21(6&7), pp.1433-1460.
 201. Zhan, X., Sohlberg, R.A., Townshend, J.R.G., DiMiceli, C., Carroll, M.L., Eastman, J.C., Hansen, M.C. and DeFries, R.S. (2002). Detection of land cover changes using MODIS 250 m data. *Remote Sensing of Environment*, Vol. 83, pp.336-350.
 202. Zhou, L.M., Tucker, C.J., Kaufmann, R.K., Slayback, D., Shabanov, N.V. and Myneni, R.B. (2001). Variations in northern vegetation activity inferred from satellite data of vegetation index during 1981 to 1999. *Journal of Geophysical Research*, Vol. 106(D17), pp.20,069-20,083.
 203. Zoran, M. and Stefan, S. (2006). Climatic changes effects on spectral vegetation indices for forested areas analysis from satellite data. In: *Proceedings of the 2nd Environmental Physics Conference*, 18-22 February 2006, Alexandria, Egypt, pp.73-83.