Volume 20 Issue 36
Originally issued as B3



Global Journal of human-social science: B
Geography, Geo-Sciences, Environmental Science & Disaster Management
Volume 20 Issue 3 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-460X & Print ISSN: 0975-587X

Significance of Baddeleyite for Paleoproterozoic PGE Deposits with Pt-Pd and Cu-Ni Reefs (North-Eastern Fennoscandian Shield): New Results of U-Pb and LA-ICP-MS Studies

By Bayanova T.B., Drogobuzhskaya S.V., Subbotin V.V., Serov P.A., Steshenko E.N., Kunakkuzin E.L. & Borisenko E.S.

Abstract- Baddeleyite is a significant mineral successfully applied in the U-Pb geochronology for the precise dating of mafic rocks from layered intrusions with the platinum group element (PGE) and Cu-Ni mineralization. The Fennoscandian Shield hosts several layered Pt-Pd, Co-Cr-Ni, and Ti-V occurrences in the Northern (Karelian) and Southern (Karelian-Finnish) belts. The aim of this study is to estimate the content and distribution of rare earth elements (REE) in baddeleyite and to calculate temperatures (T,˚C) of the U-Pb system closure and baddeleyite crystallization compared to zircon from Cu-Ni and Pt-Pd deposits in the north-eastern Fennoscandian Shield. For the first time, baddeleyite crystals from Cu-Ni (Monchepluton) and Pt-Pd (Monchetundra) reefs of the Monchegorsk ore area have been studied in situ by the laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) to measure the U-Pb age of formation and the REE content.

Keywords: baddeleyite; layered intrusions; U-Pb analysis; laser ablation inductively coupled plasma mass spectrometry; fennoscandian shield.

GJHSS-B Classification: FOR Code: 059999



Strictly as per the compliance and regulations of:



© 2020. Bayanova T.B., Drogobuzhskaya S.V., Subbotin V.V., Serov P.A., Steshenko E.N., Kunakkuzin E.L. & Borisenko E.S. This is a research/ review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Significance of Baddeleyite for Paleoproterozoic PGE Deposits with Pt-Pd and Cu-Ni Reefs (North-Eastern Fennoscandian Shield): New Results of U-Pb and LA-ICP-MS Studies

Bayanova T.B. [α], Drogobuzhskaya S.V. [σ], Subbotin V.V. [ρ], Serov P.A. [ω], Steshenko E.N. [¥], Kunakkuzin E.L. [§] & Borisenko E.S. [χ]

Abstract- **Baddeleyite is a significant mineral successfully applied in the U-Pb geochronology for the precise dating of mafic rocks from layered intrusions with the platinum group element (PGE) and Cu-Ni mineralization. The Fennoscandian Shield hosts several layered Pt-Pd, Co-Cr-Ni, and Ti-V occurrences in the Northern (Karelian) and Southern (Karelian-Finnish) belts. The aim of this study is to estimate the content and distribution of rare earth elements (REE) in baddeleyite and to calculate temperatures (T, °C) of the U-Pb system closure and baddeleyite crystallization compared to zircon from Cu-Ni and Pt-Pd deposits in the north-eastern Fennoscandian Shield. For the first time, baddeleyite crystals from Cu-Ni (Monchepluton) and Pt-Pd (Monchetundra) reefs of the Monchegorsk ore area have been studied *in situ* by the laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) to measure the U-Pb age of formation and the REE content.**

Keywords: baddeleyite; layered intrusions; U-Pb analysis; laser ablation inductively coupled plasma mass spectrometry; fennoscandian shield.

I. Introduction

Baddeleyite is a valued mineral for the U-Pb dating of PGE deposits (Bayanova, 2006; Mungall et al., 2016). Compared to zircon, it is more reliable for precise U-Pb dating of deposits in the north-eastern Fennoscandian (Baltic) Shield, since it is genetically magmatic. In contrast, zircon can be metamorphic, hydrothermal or occur as xenocrysts. The study of the trace element composition of zircon is a common practice, while geochemical characteristics of baddeleyite are poorly studied and contradictory. Thus, the value of Ce-anomaly varies, and the Eu-anomaly is absent in some analyses (Reischman et al., 1995; Zircon, 2003; Schaltegger et al., 2017). Also, recent studies limelight the crucial role of baddeleyite in the reconstruction of the supercontinents breakup in the history of the Earth's evolution (Bayanova et al., 2009; Bayanova et al., 2017; Bayanova et al., 2019 a; Bayanova et al., 2019 b; Mitrofanov et al., 2019).

In Paleoproterozoic PGE layered intrusions, baddeleyite is found in Pt-Pd and Cu-Ni deposits of the Monchegorsk ore area (Fig. 1).

Author σ: Geological Institute KSC RAS, 14 Fersman Street, 184209 Apatity, Russia. e-mail: tamarapoi@yandex.ru
Author ρ ω ¥ § χ: Geological Institute KSC RAS, 14 Fersman Street, 184209 Apatity, Russia.
Author σ: Tananaev Institute of Chemistry and Technology of Rare Elements and Mineral Raw Materials KSC RAS, Apatity, 184209, Russia.

© 2020 Global Journals



Fig. 1: Generalized geological map of the north-eastern Fennoscandian Shield and location of Paleoproterozoic layered mafic intrusions (Mitrofanov et al., 2005).

The aim of this study is (i) to estimate the REE content and distribution in baddeleyite, (ii) to calculate temperatures (T,°C) of the U-Pb system closure and baddeleyite crystallization compared to zircon from Cu-Ni and Pt-Pd deposits in the Monchegorsk ore area, hosting the recently discovered Pt-Pd Vurechuayvench deposit (north-eastern Fennoscandian Shield, Russia).

II. Resources and Techniques

Baddeleyite was extracted from gabbro-anorthosites with the Pt-Pd mineralization in the middle of the Monchetundra massif (Fig. 1). The age of baddeleyite was estimated (Nerovich et al., 2014) by the U-Pb method at 2476±5 Ma and 2471±3 Ma (Fig. 2, Table 1).



Fig. 2: Isotope U-Pb plots with Concordia for baddeleyite (*bd*) at the western slope of the Monchetundra: *a* – medium- to coarse-grained gabbronorite-anorthosite, partly amphibolite, *b* – medium- to coarse-grained leucogabbronorite.

© 2020 Global Journals

Table 1: Isotope U-Pb data on baddeleyite from the Monchetundra massif.

Sample, No.	Weighted sample, mg	Contents, ppm		Isotope composition of Pb*			Isotope ratio and age, Ma**			Rho
		Pb	U	$^{206}Pb/^{204}Pb$	$^{206}Pb/^{207}Pb$	$^{206}Pb/^{208}Pb$	$^{207}Pb/^{235}U$	$^{206}Pb/^{238}U$	$^{207}Pb/^{206}Pb$	
for baddeleyite and gabbronorite-anorthosite (medium- to coarse-grained, partly amphibolized)										
1	0.25	94.48	114.75	86	3.3003	2.2134	9.92226	0.450763	2500	0.70
2	0.20	57.60	123.14	570	5.5602	11.459	9.08165	0.417879	2430	0.93
3	0.25	30.09	67.86	557	5.6496	8.0718	8.19067	0.385383	2392	0.88
for baddeleyite from medium- to coarse-grained gabbronorite-anorthosite										
1	0.50	110.70	244.90	1478	5.9187	23.690	9.504470	0.429716	2460	0.94
2	0.35	152.60	359.10	3510	6.1640	35.011	9.119096	0.413367	2441	0.95
3	0.50	60.479	136.81	830	5.5615	13.663	8.820570	0.400447	2453	0.91
4	0.20	98.025	246.35	1539	6.3461	24.580	8.368770	0.382377	2437	0.83

** The ratios are corrected for blanks of 0.8 ng for Pb and 0.04 ng for U and mass discrimination 0.12 ± 0.04%.*
*** Correction for common Pb is estimated for the age, according to Stacey and Kramers (1975).*

The REE content and distribution in baddeleyite were estimated by the following technique. The method of the electron (LEO-1415) and optic (LEICA OM 2500 P, camera DFC 290) spectroscopy was used to study the morphology of samples. Points for local analyses on baddeleyite crystals were selected based on the study of their back-scattered electron (BSE) and cathodoluminescence (CL) images.

Contents of REE and other elements were estimated *in situ* by ICP-MS on an ELAN 9000 DRC-e (Perkin Elmer) quadrupole mass spectrometer equipped with a 266 nm UP-266 MACRO laser (New Wave Research). LA-ICP-MS was performed using argon with a repetition rate of 10 Hz, pulse duration of 4 ns, the energy density of 14-15 J/cm^2 at a spot with a diameter of 35-100 μm or using scanning "in a line" (length 35-70 μm), monitoring and measuring produced craters. NIST 612 glass with the known concentrations of REE, U, Ti, and Th of 40 ppm was used for external calibration as a multi-point calibration forced through the origin after blank correction (Pearce et al., 1997, Certificate of Analysis, 2012). NIST SRM 610 sample (450 ppm concentration) was used to check the accuracy of estimations (Yuan et al., 2004; Jochum et al., 2011). The laser beam diameter was changed, while the rest parameters were stable: from 35 to 240 μm (point sampling) and from 20 to 155 μm (r = 0.999) (scanning "in a line"). As for calibration standards, measurements of the elements were in the range of 15% relative deviations. Determination limits were within 0.01 ppm, a diameter of the laser beam of 155 μm. It complies with the available data (Yuan et al., 2004). This technique was tested, using analyses of internationally approved standard zircon samples 91500, TEMORA 1, Mud Tank, and inter-laboratory cross-checks (Boynton, 1984).

III. Results

Table 2 and Figure 3 provide new data on the contents of REE and other elements in baddeleyite from Pt-Pd occurrences of the Monchetundra massif.

Baddeleyite from vein pegmatites with the gabbronorite composition from the Monchepluton with Cu-Ni reefs (Mt. Nyud, Terrace deposit) was studied. Its U-Pb age was estimated at 2505±5 Ma (Bayanova, 2006). Figures 4 and 5 display new LA-ICP-MS data on baddeleyite, which was measured along and across its section.

© 2020 Global Journals



Fig. 3: Chondrite-normalized plots of REE distribution in baddeleyite from medium- to coarse-grained leucogabbronorite of Pt-Pd occurrences, Monchetundra massif (Boynton et al., 1984).



Fig. 4: Chondrite-normalized plots of REE distribution in baddeleyite from gabbronorite-anorthosite of Pt-Pd occurrences, Monchetundra massif (Boynton et al., 1984).

© 2020 Global Journals



Fig. 5: Chondrite-normalized plots of REE distribution in baddeleyite from vein pegmatites of the gabbronorite composition, Cu-Ni Terrace deposit, Mt. Nyud, Monchetundra massif (Boynton et al., 1984).

Notably, spectra of REE distribution in baddeleyite grains from PGE rocks of the Monchegorsk ore area indicate low concentrations of light elements, high concentrations of heavy ones, steep positive slope with a negative Eu-anomaly and a positive Ce-anomaly. The spectra also reflect the magmatic origin of baddeleyite (Zircon, 2003; Watson et al., 2006).

IV. Conclusion

For the first time, the provided research revealed a direct relation between the REE content in baddeleyite and the formation of Pt-Pd and Cu-Ni reefs. The higher concentrations of HREE in baddeleyite, the higher temperatures of the U-Pb systematics closure and formation are. Pt-Pd reefs are likely to occur under such conditions. Cu-Ni reefs form at lower temperatures of the U-Pb systematics closure and crystallization of accessory minerals. These occurrences display low HREE and a wide range of LREE concentrations (Table 2).

V. Funding

The research has been funded by grants of the Russian Foundation for Basic Research, 18-35-00246, 18-35-00152, 18-05-70082 (Arctic Resources) and Presidium RAS Program No. 1.8. State Order research contracts No. 0230-2018-0002 and 0231-2015-0005.

Acknowledgments

This paper is devoted to the memory of outstanding geologists Academicians F.P. Mitrofanov and V.T. Kalinnikov, geochemists E.B. Bibikova and J. Wasserbourg, who kindly submitted an artificial tracer 205 to create a new U-Pb spike for precise U-Pb dating of baddeleyite, in particular.

References Références Referencias

1. Mungall, J., Kamo, S.L. and McQuade, S. (2016) U-Pb Geochronology Documents Out-of-Sequence Emplacement of Ultramafic Layers in the Bushveld Igneous Complex of South Africa. Nat. Comm. 7. https://doi.org/10.1038/ncomms1338
2. Bayanova T.B. (2006) Baddeleyite: A Promising Geochronometer for Alkaline and Basic Magmatism // Petrol. 14 (2). 187-200.
3. Reischmann, T.; Brugmann, G.E.; Jochum, K.P.; Todt, W.A. (1995) Trace element and isotopic composition of baddeleyite. Mineral Petrology, 53, 155-164.
4. Zircon. Rev. Miner. Geochem. (2003) (Eds. John M. Hanchar & Paul W.O. Hoskin). 53. 500.
5. Nerovich L.I., Bayanova T.B., Serov P.A. and Elizarov D.V. (2014) Magmatic Sources of Dikes and Veins in the Moncha Tundra Massif, Baltic Shield: Isotopic-Geochronologic and Geochemical Evidence// Geochem. Intern. 52 (7). 548–566.
6. Pearce, N.J.G.; Perkins, W.T.; Westgate, J.A. (1997) A compilation of new and published major and trace element data for NIST SRM 610 and NIST SRM 612 glass reference materials. Geostand. Newslett. 21(1), 115-144.
7. Certificate of Analysis. Standard Reference Material 612. Trace Element in Glass (2012) // NIST.1-4.
8. Jochum K.P., Weis U., Stoll B., Kuzmin D., Yang Q., Raczek I., Jacob D.E., Stracke A., Birbaum K., Frick D.A., Günther D., and Enzweiler J., (2011) Determination of Reference Values for NIST SRM

© 2020 Global Journals

610-617 Glasses Following ISO Guidelines. Geostandards and Geoanalytical Research. 35. 4. 397-429.
9. Yuan H., Gao S., Liu X., Li H., Günther D., Wu F., (2004) Accurate U–Pb age and trace element determinations of zircon by laser ablation inductively coupled plasma-mass spectrometry. Geostand. Geoanal. Res. 28. 353–370.
10. Nikolaev, A.I.; Drogobuzhskaya, S.V.; Bayanova, T.B., Kaulina, T.V.; Lyalina, L.M.; Novikov, A.I.; Steshenko, E.N. (2016) REE distribution in zircon from reference rocks of the Arctic region: evidence from study by the LA-ICP-MS method. Doklady Earth Sci. 470(2), 1037-1041.
11. Boynton, W.V. (1984) Cosmochemistry of the rare Earth elements: Meteorite Studies. In: Henserson, P., Ed., Rare Earth Element Geochem. Elsevier, Amsterdam, 63-114. http://dx.doi.org/10.1016/B978-0-444-42148-7.50008-3
12. Schaltegger U., Davies J.H.F.L. (2017) Petrochronology of zircon and baddeleyite in igneous rocks: Reconstructing magmatic processes at high temporal resolution//Rev. Mineral. Geochem.83 (1). 297-328.
13. Watson, E.B., Wark, D.A. and Thomas, J.B. (2006) Crystallization Thermometers for Zircon and Rutile. Contrib. Mineral. Petrol. 151. 413-433. https://doi.org/10.1007/s00410-006-0068-5
14. Stacey, J.S.; Kramers J.D. (1975) Approximation of terrestrial lead isotope evolution by a two-stage model. In Earth and Planet. Sci. Lett. 26(2). 207-221.
15. Mitrofanov, F.P. (2005) New types of mineral raw materials in the Kola Province: discoveries and perspectives. Smirnovsky Collection of works. Moscow: VINITI RAS. pp. 39 – 53.
16. Heaman, L.M.; Le Cheminant, A.N. (1993) Paragenesis and U-Pb systematics of baddeleyite (ZrO). Chem. Geol. 110, 95-126.
17. Bayanova T., Ludden J. & Mitrofanov F. Timing and duration of Palaeoproterozoic events producing ore-bearing layered intrusions of the Baltic Shield: metallogenic, petrological and geodynamic implications / In: Reddy S.M., Mazumder R., Evans D.A.D. & Collins A.S. (eds) // Palaeoproterozoic Supercontinents and Global Evolution. Geological Society, London, Special Publications, 2009. N. 323. P. 165-198.
18. Bayanova T., Mitrofanov F., Serov P., Nerovich L., Yekimova N., Nitkina E. and Kamensky I. Layered PGE Paleoproterozoic (LIP) Intrusions in the N-E Part of the Fennoscandian Shield – Isotope Nd-Sr and $^{3}He/^{4}He$ Data, Summarizing U-Pb Ages (on Baddeleyite and Zircon), Sm-Nd Data (on Rock-Forming and Sulphide Minerals), Duration and Mineralization/Geochronology – Methods and Case Studies/Edited by Nils-Axel Mörner//INTECH, 2014. P. 143-193.
19. Mitrofanov F. P., Bayanova T. B., Ludden J. N., Korchagin A. U., Chashchin V. V., Nerovich L. I., Serov P. A., Mitrofanov A. F., Zhirov D. V.. Origin and Exploration of the Kola PGE-bearing Province: New Constraints from Geochronology/Ore Deposits: Origin, Exploration, and Exploitation/Edited by Sophie Decree and Laurence Robb//Geophysical Monograph Series. Wiley. 2019. P. 3-36. Print ISBN: 9781119290537 |Online ISBN: 9781119290544 |DOI:10.1002/9781119290544.ch1
20. Bayanova T., Korchagin A., Mitrofanov A., Serov P., Ekimova N., Nitkina E., Kamensky I., Elizarov D., Huber M. Long-lived mantle plume and polyphase evolution of Palaeoproterozoic PGE intrusions in the Fennoscandian Shield//Minerals. – 2019. – № 9 (1). – V. 59. doi:10.3390/min9010059
21. Bayanova T., Subbotin V., Drogobuzhskaya S., Nikolaev A., Steshenko E. Baddeleyite from Large Complex Deposits: Significance for Archean-Paleozoic Plume Processes in the Arctic Region (NE Fennoscandian Shield) Based on U-Pb (ID-TIMS) and LA-ICP-MS Data // Open Journal of Geology. – 2019. – V. 9. – No 8. – P.474-496. DOI: 10.4236/ojg.2019.98031

© 2020 Global Journals

Table 2: LA-ICP-MS data on contents of REE and other elements in grains of baddeleyite and zircon from Pt-Pd and Cu-Ni reefs of the Monchegorsk ore area.

Contents (average in crystal), *ppm*																						
Sample			Ti	Y	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	Hf	Th	U	Σ REE	T°C
Sample *1*	*Bd*	2	645.2	230. 3	0.75	5.65	0.29	2.90	1.52	0.18	5.23	1.51	20.5	8.90	46.8	11.4	107.4	25.5	461 7	2.82	68.2	238.5	> 1000° C
		3	468. 7	429.5	12.1	29.4	2.96	16.9	7.08	0.21	18.9	4.72	50.9	17.8	83.1	18.8	162.2	32.7	297 0	9.51	212.6	457.8	
		7	1212	848. 5	7.89	39.0	4.19	20.1	10.3	0.41	18.0	4.83	63.8	25.7	155.5	34.9	315.3	72.0	7453	73.8	172.4	771.9	
Sample *2*	*Bd*	4	260.0	108.6	0.38	2.11	0.43	1.60	1.58	0.55	3.46	0.88	11.1	4.80	24.6	8.30	69. 2	14.5	306 0	12.6	58.2	143.5	
		7-1-2	432.2	264. 0	1.54	5.35	0.41	2.50	2.45	0.18	5.47	1.94	26.2	9.30	50.1	10.5	94.6	18.1	1413	25.9	89.4	228.6	
		7-3-4	549. 9	335.9	1.96	6.80	0.53	3.17	3.11	0.22	6.96	2.47	33.4	11.8	63.8	13.4	120.4	23.0	1797	32.9	113.7	291.0	
		9	878.5	155.7	1.95	12.9	1.37	11.5	4.23	0.30	4.94	0.62	11.3	5.85	30.3	5.85	67.1	12.4	5653	31.1	108.1	170.6	
Sample *3*	*Bd*	I 2	1637	21.1	0.09	1.76	0.06	0.41	0.18	0.09	0.54	0.16	1.45	0.69	3.41	0.71	7.22	1.33	6132	2.83	196.5	18.1	
		II 3	2358	38.6	0.01	3.95	0.01	0.06	0.01	0.02	1.00	0.06	2.49	1.82	6.49	1.48	16.6	3.60	5761	3.04	213.8	37.6	
		II 2	3949	132.9	3.88	26.1	0.60	5.96	1.52	2.22	5.58	1.40	14.4	4.19	19.1	4.81	49.1	10.6	10225	27.6	1057	149.5	
		II 1	940. 1	20.4	0.32	1.56	0.03	0.23	0.13	0.07	0.51	0.19	1.96	0.92	3.90	0.77	6.47	2.16	3068	3.51	135.8	19.2	
		2-1	6015	52.4	0.20	4.33	0.03	0.57	0.81	0.03	1.12	0.69	6.06	2.34	11.6	2.56	28.2	3.96	17905	13.5	306.1	62.5	
		1-1	3742	60.6	0.17	4.56	0.04	0.24	0.37	0.14	1.35	0.62	7.18	2.56	12.5	3.04	29.3	4.85	809 6	14.7	185. 8	66.9	
		1-3	4876	108.1	0.60	7.16	0.14	1.20	0.69	0.11	2.81	1.14	10.7	4.84	23.4	6.16	59. 8	9.5 9	175 43	10.6	303. 3	128.3	
	Zr	II 1	51.1	638. 3	0.18	15.6	0.31	5.51	7.77	0.80	32.6	6.63	71.7	20.9	75.5	12.5	96. 6	14.2	900 9	559. 0	242. 5	360.8	aver. 894° C
		II 2	44.6	322. 1	0.13	15.7	0.95	7.47	8.17	1.07	26.7	5.58	43.2	10.4	32.4	4.96	32. 4	3.99	731 1	166. 6	71.9	193.1	
		II 3	35.2	718. 0	0.03	13.4	0.06	0.70	2.03	0.14	15.3	4.66	61.2	22.2	100. 0	18.3	148.5	23.1	913 3	207. 9	121. 9	409.6	
		II 4	37.6	392. 1	0.34	11.9	0.30	3.17	4.54	0.80	16.9	4.16	42.3	11.9	44.3	7.60	60. 7	8.91	8570	346. 2	217.6	217.8	
		III 1	13.1	649. 3	0.22	14.3	0.43	6.13	7.57	0.38	30.8	7.44	74.8	21.6	71.9	11.4	90. 2	10. 7	369 6	325.8	122. 5	347.9	
		III 2	64.1	3112	0.26	69.2	1.43	17.4	31.7	2.45	140. 0	32.0	327. 4	100. 1	386. 3	63.1	524	73.9	837 4	127 1	428.6	1769	
		III 3	118. 4	703. 4	0.13	19.9	0.85	11.2	10.0	2.03	34.7	8.69	81.8	21.9	71.8	13.0	104.6	13. 5	214 22	166 9	664.8	394.1	

Note: 1- baddeleyite from gabbronorite -anorthosite, 2 - baddeleyite from medium- to coarse -grained leucogabbronorite, 3- sample from vein pegmatites of the gabbronorite composition. Temperature of zircon and baddeleyite crystallization is calculated according to [3].

© 2020 Global Journals