

# Changes in Earth Cover and Relations with the Equation Universal Soil Losses in Basin of the Western Plateau of Sao Paulo, Brazil

Alyson Bueno Francisco

Received: 11 June 2021 Accepted: 2 July 2021 Published: 15 July 2021

## Abstract

The expansion of sugarcane crop in watersheds, located in the regions of Marilia and Presidente Prudente, in the interior of São Paulo, by replacing degraded pastures, is altering soil cover and influencing erosive processes. This article aims to analyze the land cover changes in the basin of Confusion's stream and relate to the parameters of the factors of the Universal Soil Loss Equation. The land cover charts were generated with images of CBERS's satellite dated 2009 and 2018 by supervised classification. The areas of the land cover classes were related to factor C of the USLE and data estimates of secondary sources of erosivity factor (R), erodibility factor (K) and topographic factor with the data of slope (LS). From these parameters, an estimate of soil loss was presented for the basin of Confusion's stream. The soil loss in the basin of Confusion's stream was estimated by 2018 data at 2,484 t/ha/y.

**Index terms**— hydrographic basin; erosion; cartography; GIS.

## 1 Introduction

he first estimates of soil losses in Brazil were analyzed in the 1940s using experimental plots at the Agronomic Institute of Campinas (BERTONI; LOMBARDI ??ETO, 1999).

3 The Universal Equation of Soil Losses was developed from the studies of Wischmeier and Smith ??1961 apud  
4 LAFLEN;MOLDENHAUER, 2003) from 10,000 data on soil loss rates in experimental plots in the United States  
5 of America.

6 Soil losses in Brazil are estimated at approximately 848 million tons per year (MERTEN; MINELLA, 2013).  
7 Fast-paced soil losses in the degraded areas of the Western Plateau of São Paulo represent environmental and  
8 economic damage.

The scale of watersheds has repercussions in recent years due to the details of the guaranteed by advances in geographic technologies. On the local scale, the particularities of each rural producer are found and mitigating measures are applied in the face of land degradation conditions. The dialogue with rural producers takes place through the technical knowledge of public agencies and researchers guide these agencies through extension projects. The definition of the concepts and principles of water management is essential to consolidate the necessary measures in river basins. Ideals based on rational water use are important to guide political decisions and favor of soil conservation measures and recovery of degraded areas.

6 The geoprocessing techniques used met the research needs, confirming that its use improves the processes and  
7 phases of spatial analysis. In addition, the created database can be replicated, corrected and updated at any time,  
8 which makes it dynamic and applicable to the most diverse spatial representation demands of the information  
9 contained therein (PIROLI, 2013).

The watershed is a territorial and physical unit present in nature bounded by drainage. Nature presents its diversity and researchers need to avoid generalizing small scales and seek to understand natural phenomena in field research.

In this sense, fieldwork is necessary, as the main methodology of the geographer, to investigate the nature of each hydrographic basin and present the diagnoses (inventories), to carry out environmental planning from the

## 4 RESULTS AND DISCUSSION

---

45 perspective of water resources management with adequate prognoses to ensure the future availability of drinking  
46 potable water (GUIMARÃES, 1999).

47 From the development of Geographic Information Systems, spatial data from watersheds served as parameters  
48 applying the Universal Equation of Soil Losses, mainly by the formation of numerical elevation models. The  
49 development of erosive processes in the large hydrographic basins of tropical environments made it propitious to  
50 apply this empirical model in river basins to generate databases in Geographic Information Systems (PARVEEN;  
51 KUMAR, 2012).

52 In recent years, research on river basins and geoprocessing techniques has become more present in geographical  
53 studies. The spatiality of erosive processes involves the understanding of the aspects of hydrographic basins and  
54 the transformations that occur in rural landscapes of changes in agricultural activities. The use of conservation  
55 practices for the recovery of areas with degraded soils can contribute to the mappings performed by geographers  
56 with support in geographic information systems. Despite deforestation for cotton planting until the 1970s and  
57 later the domain of pasture planting for cattle herd farming, the Confusion Stream's basin has preserved fragments  
58 of native forest (approximately 8% of the area), including a fragment of approximately 3,000 hectares.

59 In the last 15 years, installing a sugar-alcohol plant in the southwestern part of the basin favored the increase  
60 of sugarcane planting areas (23% to 34% of the area) with the existence of terraced plantations, replacing the  
61 degraded pasture areas.

62 The clinographic letter of slope was elaborated based on data from the digital elevation model from the SRTM  
63 mission, in the Slope routine of the TerrSet GIS. The slope data served as the basis for the elaboration of the  
64 estimation of the LS factor, ramp length (slopes) and unevenness (slopes).

65 The preparation of the land cover charts were carried out in the TerrSet Geographic Information System  
66 with orbital images of the CBERS satellite dated October 2009 and February 2018, through the supervised  
67 classification routine. The geographic referencing of the images was performed at GIS Idrisi. After geographic  
68 referencing, the false-color composition was applied with bands 2, 3 and 4.

69 Using the Digitize routine, reliable samples were selected with the vectorization of polygons, whose each class  
70 to be represented had a value. With the use of the Make-sig routine, signatures are created and then the classifier  
71 is defined with maximum likelihood with the use of the Max-like routine. Then, mode filtering was applied using  
72 a size 7x7. To cut out the area of the

## 73 2 METHODOLOGY

74 Topography Mission (SRTM) data were delimited using SRTM data in the Global Mapper software. The vector  
75 that delimits the watershed was exported in Shapefile format, having as reference system the horizontal datum  
76 SIRGAS-2000.

77 Regarding the factors attributed to the Universal Equation of Soil Losses, the erodibility indexes presented by  
78 Freire et al. (1992). The rainfall erosivity (R) was estimated based on the reference of Francisco (2017) when  
79 presenting the values for an adjacent basin 15 km away from the Confusion Stream's basin. The types of soils  
80 identified in the hydrographic basin were adopted by the map presented by Oliveira et al. (1999). The C factor  
81 of soil management forms, by the land cover classes, followed the reference adopted by Pinto (1991). The factor  
82 conservation practices (P) was estimated based on the indices of calculations expressed by Bertoni and Lombardi  
83 Neto (1999). Table 2 shows the relationship between the areas of the land cover classes and the C factor of land  
84 use applied in the Universal Equation of Soil Losses. The data show factor C from 810.17 to 994.31, an increase  
85 of 18.5% in erosive potential according to land cover changes in 9 years.

## 86 3 III.

## 87 4 Results and Discussion

88 The table 3 presents the relation of the areas of the land cover classes with the predominant types of soils by  
89 the K factor of erodibility. The erodibility factor remained practically stable in 9 years, being present the addition  
90 of native forest areas present in hydromorphic soils of primary forests and a decrease in exposed soil areas, very  
91 conducive to the development of laminar erosion. When considering the average length of 100 m of ramp in the  
92 basin and the mean slope of 5.67% or 10°, the topographic factor was estimated at 0.006.

93 The factor of conservation practices was calculated based on the mean slope of 10°, being obtained in 0.19342.

94 When considering the estimated factors of the USLE for the basin, the rates of soil losses were estimated in  
95 2009 by the land cover classes in table 5. The Confusion Stream's basin, with 46,760 ha, presents an estimate of  
96 soil losses at 1,909.24 t/ha/year in 2009.

97 When considering the estimated factors of the USLE for the basin, the rates of soil losses were estimated  
98 in 2018 by the land cover classes in table 6. The Confusion Stream's basin, with 46,760 hectares, presents an  
99 estimate of soil losses at 2,484.08 t/ha/year in 2018.

100 IV.

## 101 5 Conclusions

102 The estimates of soil losses are important for the management of watersheds with the application of mapping in  
103 the identification of the most degraded areas susceptible to erosion. The mapping of the current conditions of the  
104 hydrographic basin (inventory) favors the diagnosis of degradation conditions and proposals for environmental  
105 recovery.

106 Brazil is a country with large territorial extensions and watersheds that are distributed in several municipalities  
107 and states and it is important to implement policies for the management of water resources through partners  
108 with the various territorial units. Brazilian municipalities need to draw up an inventory with maps and databases  
109 of watersheds to favor water management.

110 Soil conservation is an interdisciplinary theme and can integrate several professionals from the technical and  
111 research fields to support public agencies in the proposals to recover areas degraded by rigorous erosive processes.

112 Soil degradation conditions require the improvement of more empirical scientific techniques and with the  
113 support of geographic information systems to contribute to the monitoring of erosive processes. <sup>1</sup>




Figure 1: The

113

<sup>1</sup>© 2021 Global Journals Year 2021 B Changes in Earth Cover and Relations with the Equation Universal Soil Losses in Basin of the Western Plateau of São Paulo, Brazil

## 5 CONCLUSIONS




Figure 2: Figure 1 :)Figure 2 :




Figure 3: B




Figure 4: Figure 3

1

Land covers classes	Area in 2009 (ha)	%	Area in 2018 (ha)	%
Urban areas and roads	0.58	0.01	1.55	0.003
Native forest	3,610.00	7.72	3,913.28	8.37
Water channel	1.42	0.01	1.58	0.003
Agricultural crop	10,679.00	22.84	15,768.36	33.72
Pasture	23,085.00	49.75	22,571.32	48.27
Land wood	326.00	0.70	-	-
Bare soil	8,698.00	18.75	3,504.06	09.61

Figure 5: Table 1 :

2

Land covers classes	Factor C	Area in 2009 (ha)	C (2009)	Area in 2018 (ha)	C (2018)
Native forest	0.0004	3,610	0.00003	3,913.28	0.00003
Agricultural crop	0.0500	10,679	0.11420	15,768.36	0.16860
Pasture	0.0075	23,085	0.37000	22,571.32	0.36200
Land wood	0.0450	326	0.00700	-	-
Bare soil	0.1000	8,698	0.18601	3,504.06	0.00370

Figure 6: Table 2 :

## 5 CONCLUSIONS

---

3

Classe de cobertura da terra	Soil kind	Factor K	Área in 2009 (ha)	K (2009)	Área in 2018 (ha)	K (2018)
Native forest	Hydric soil	0.003	3,610	0.00023	3,913.28	0.00025
Agricultural crop	Clay soil	0.051	10,679	0.01165	15,768.36	0.01719
Pasture	Oxisoil	0.016	23,085	0.00799	22,571.32	0.00772
Land wood	Oxisoil	0.016	326	0.00011	-	-
Bare soil	Clay soil	0.051	8,698	0.00949	3,504.06	0.00382

Figure 7: Table 3 :

61  
(

Figure 8: )

4

Classe de cobertura da terra	Área in 2009 (ha)	R (2009)	Área in 2018 (ha)	R (2018)
Native forest	3,610	564.66	3,913.28	610.92
Agricultural crop	10,679	1,667.16	15,768.36	2,461.64
Pasture	23,085	3,603.94	22,571.32	3,523.7
Land wood	326	50.89	-	-
Bare soil	8,698	1,357.9	3,504.06	547.03
?		7,244.55		7,143.29

Figure 9: Table 4 :

5

Land covers	Factor R	Factor K	Factor LS	Factor C	Factor P	Ei	A (t/ha/year)
Native forest	564.66	0.00023	0.006	0.00003	0.19342	2.10 -9	0.0002
Agricultural crop	1,667.16	0.01165	0.006	0.11420	0.19342	0.0257	1,201.73
Pasture	3,603.94	0.00799	0.006	0.37000	0.19342	0.0123	577.51
Bare soil	1,357.9	0.00949	0.006	0.18601	0.19342	0.0027	130

Figure 10: Table 5 :

---

**6**

Land covers	Factor R	Factor K	Factor LS	Factor C	Factor P	Ei	A (t/ha/year)
Native forest	610.92	0.00023	0.006	0.00003	0.19342	2.10 -9	0.0002
Agricultural crop	2,461.64	0.01165	0.006	0.11420	0.19342	0.038	1,777.23
Pasture	3,523.7	0.00799	0.006	0.37000	0.19342	0.012	566
Bare soil	547.03	0.00949	0.006	0.18601	0.19342	0.003	140.85

Figure 11: Table 6 :

## **5 CONCLUSIONS**

---

- 
- 114 [Freire et al. ()] 'Erodibilidade de alguns solos do oeste do estado de São Paulo'. O Freire , M C T F Godoy , W  
115 A Carvalho . *Revista de Geografia* 1992. 1 (11) p. .
- 116 [Perusi et al. ()] 'Erosividade das chuvas no extremo oeste do Estado de São Paulo nos anos de 1998 e 1999'. M  
117 C Perusi , V Zero , J T G Tommaselli , N Brigatti . *Caderno Prudentino de Geografia* 2004. 26 (12) p. .
- 118 [Piroli ()] *Geoprocessamento aplicado ao estudo do uso da terra das áreas de preservação permanente dos corpos*  
119 *d'água da Bacia Hidrográfica do Rio Pardo. Thesis (Geoprocessing and Remote Sensing)*, E L Piroli . 2013.  
120 p. 150. São Paulo State University
- 121 [Parveen and Kumar ()] 'Integrated Approach of Universal Soil Loss Equation (USLE) and Geographical  
122 Information System (GIS) for Soil Loss Risk Assessment in Upper South Koel Basin'. R Parveen , U Kumar  
123 . *Jharkhand. Journal of Geographic Information System* 2012. 6 (4) p. .
- 124 [Oliveira et al. ()] *Map of Soils in the State of São Paulo*, J B Oliveira , M N Camargo , M ; Rossi , Calderano  
125 , B Filho . Scale: 1:500.000. 1999. Agronomic Institute of Campinas.
- 126 [Pinto ()] 'Sensoriamento Remoto e integração de dados aplicados no estudo da erosão dos solos: contribuição  
127 metodológica'. S A F Pinto . *Thesis (Physical Geography)* 1991. p. 185. Universidade de São Paulo
- 128 [Merten and Minella ()] *The expansion of Brazilian agriculture: soil erosion scenarios. International Soil and*  
129 *Water Conservation Research*, G H Merten , J P G Minella . 2013. p. .
- 130 [Laflen and Moldenhauer ()] *The Story USLE: pioneering soil erosion prediction*, J M Laflen , W C Moldenhauer  
131 . 2003. Beijing, China: World Association of Soil and Water Conservation.
- 132 [Guimarães ()] *Trabalhos de campo em bacias hidrográficas: os caminhos de uma experiência em Educação*  
133 *Ambiental*, E M A Guimarães . 1999. p. 184. Universidade Estadual de Campinas (Dissertation (Master in  
134 Geography)