

Health and Nutritional Status of a Primitive Tribe of Madhya Pradesh: Bhumia

Ajeet Jaiswal¹

¹ Pondicherry University, Puducherry, India

Received: 8 December 2012 Accepted: 31 December 2012 Published: 15 January 2013

Abstract

Health and nutritional status of an individual depends on the food he eats. Man needs a wide range of nutrients to lead a healthy and active life and these are derived through the diet they consume daily. Good nutrition is a basic component of health. This paper deals with the utility of various anthropometric cut-off points in the evaluation of nutritional Health status. Total 616 subjects studied from 201 household of the five villages of Baigachak area. Anthropometric measurements taken were height and weight. Dietary data was collected using 24hr recall method. The extent of malnutrition for preschool children was assessed by SD classification and the nutritional status of adults was assessed by BMI classification.

Index terms— bhumia tribe, nutritional status, nutrient intake.

1 Introduction

Madhya Pradesh is the largest tribal state, with 23% of the total tribal population in the country & of 46 tribal groups in Madhya Pradesh six are considered to be primitive, based on neglect in the past, backwardness and low economic level. The Baiga, Bhumia tribe one of these six primitive tribes are mainly in five districts (Madla, Shahdol, Bilaspur, Bhalaghat and Rajnandgaon). Bhumia tribes are also located in the states of Bihar, Madhya Pradesh, Maharashtra, Orissa, and West Bengal. According to 1961 Census Data, Bhumia was Author : Pondicherry University, Puducherry, India. e-mail: rpgajeet@gmail.com reported from 9 districts of Madhya Pradesh totaling to 54520 (M.L. ??atel 2007). Baigas or Bhumia are one of the oldest aboriginal tribes and classified as one of the primitive tribe of Madhya Pradesh on the basis of preagricultural technology, low literacy and stagnant and diminishing population (Tewari 1984). Shifting cultivation depicted as an important means of livelihood.

Tribal peoples are acknowledged to have very close association with the ecosystem and the environment because of their fulfillment of daily nutritional requirements with food foraged from nature. Inadequacies in nutritional intake or under-nutrition can be considered a major source of many adverse effects on the growth and health of individuals (Gordon et al., 1968). Knowledge of the nutritional status of a community or a region is necessary to have a comprehensive idea about its development process, as under nutrition is one of the major health problems in developing countries. It is reported that the basic cause of under nutrition and infection of people are poverty, poor hygienic condition and little access to preventive health care (Mitra 1985 ?? WHO 1990). Nutritional status can be assessed by dietary, anthropometric, biochemical and clinical methods. Ideally, a combination of methods should be used when assessing nutritional status using standardised techniques.

The Bhumia's economy is still highly depended on agricultural pursuits and collection of minor forest produces. Maize, Vargu and Rice is the staple grain and forms an important item of daily diet. The baiga people also kodo, kutki, drink pej, eat little flour and are normally content with what little that they get. One of the prime foods is pej that can be made from grounding macca or from the water left from boiling rice. Also, they eat several items from the forest that includes primarily Chirota Bhaji, Gular leaves such as Chirota, chinch, chakora, sarroota, peepal etc. They also eat BirarKand, Kadukand and other rhizomes. Mushroom is also a delicacy. Numerous fruits such as mango, char, jamun, tendu are also eaten. They hunt as well, primarily fish and small

mammals (K. Sharma 2007) Out of 35 States (28 states and 7 union territories) in India, some are identified as demographically lagging behind and Bhumia Tribes of Madhya Pradesh also falls in this category and needs a situational analysis related to nutritional and Health condition. The present paper deals with the health Abstract-Health and nutritional status of an individual depends on the food he eats. Man needs a wide range of nutrients to lead a healthy and active life and these are derived through the diet they consume daily. Good nutrition is a basic component of health. This paper deals with the utility of various anthropometric cut-off points in the evaluation of nutritional & Health status. The tribal population of the country, as per the 2001 census, is 8.43 crore, constituting 8.2% of the total population with 91.7% of them living in rural areas and 8.3% in urban areas. The population of tribes had grown at the growth rate of 24.45% during 1991-2001. More than half of the Scheduled Tribe population is concentrated in the States of Madhya Pradesh (14.51%).

T II.

2 Material and Methods

This study is a review work, done to understand the prevalence of under nutrition & health status among Bhumia tribe of Baigachak area of Dindori district in Madhya Pradesh. The implications of the discussed research will help in formulating better recommendations for further research so as to reduce this nutritional burden. The Baigachak area is spread out in 39 villages in three blocks. Total five villages were selected purposely from all three blocks for this study considering its accessibility during survey.

A total of 201 households comprising of 616 which 317 male, 299 female including 125 preschool children were randomly selected during the study. Thus the present investigation has been conducted by random sampling using pre tested, structured interview schedule. Simultaneously, group discussion and informal interview methods have been used. Observation has been conducted through semi-participant method. Anthropometric measurements were taken using standard procedure (Weiner and Lourie (1981; ?eliff, 1966). BMI was computed using the following standard equation: $BMI = \text{weight (kg)} / \text{height}^2 \text{ (m}^2\text{)}$. Nutritional status was evaluated using internationally accepted BMI guidelines using James' classification (James 1988). Body weight was measured using lever actuated balance in Kg with accuracy of 100gm with minimum clothing. Height was measured by anthropometry rod (SECA, Yogul and Halke, GmbH and Co. Hamburg, Germany). Diet survey was carried out in every fifth household using 24 hours Dietary recall method (Thimmayama 1987). The nutrient intake was calculated using food tables for Indian foods (Gopalan et al. 1990) and food intakes were compared with the balanced diets recommended for Indians (ICMR 1981). The intake of nutrients was compared with the recommended Dietary allowances for Indians (ICMR 1990). The results were compared with the tribal data of National Nutrition Monitoring Bureau (NNMB, 2009).

The data analysis was carried out using statistical software package SPSS 13.0 version. Univariate analysis using t-test was applied to evaluate the statistical significance. Mean and Standard Deviation of the anthropometric data was calculated for each age group and compared with NCHS (National Center of Health statistics) standards (NCHS 1976). (NCHS data are of whole year).

3 III.

4 Results

A total of six hundred and sixteen individuals' height and weight were recorded (Table 1). Preschool girls were slightly taller than the boys of same age group. However mean weight was comparable in both genders in the same age group. In the 7-10 years age group there was no difference in the mean height of boys and girls. However adolescent girls up to 15 years were taller by about 2-3 cm and heavier by 1-2 kg as compared to boys of the same age group.

In contrast boys of 16 years old were taller by about 6-10 cm and heavier by 3-5 kg as compared to the girls of the same age group. Similar observation was made with respect to adults also. The tribals were shorter and lighter when compared with the NCHS standards (Table 2). The overall stunting was 42.2% and wasting was 36.2% and severely stunting and wasting was 20.1% and 9% respectively. Prevalence of chronic energy deficiency (BMI<18.5) through body mass index was about 77 per cent among adult population and adults females are slightly better nourished (26.0%) as compared to males (21.4%) (Table 4). Two hundred and thirty six individuals were assessed for Dietary intake information. Maize and Rice formed the bulk of Bhumias diet. The mean intake of cereals (475g/day) was higher than the recommended level ($P<0.05$).

However the intake of foodstuffs, such as pulses, green leafy vegetables, root and tubers, oil and fat, sugar and jaggery was significantly lower than recommended level (Table 5) ($P<0.05$). The milk intake (5.5 g/day) was almost negligible in Bhumias. The intake of all nutrients except calcium was significantly lower than recommended level (Table 6) ($P<0.05$).

IV.

5 Discussion

Health is a universally cherished goal. Health cannot be forced upon the people. It is a positive attribute of life and the organization of health services to all people is considered to be the key step towards development

(Srinivasan 1987). Health care is one of the most important of all human endeavours to improve the quality of life especially of the tribal people (Balgir, 1995(Balgir, , 2000(Balgir, , 2005(Balgir, & 2007)).

It implies the provision of conditions for normal, physical and mental development and functioning of human being individually as well as in a group. Health problems and health practices of tribal communities have been profoundly influenced by the inter-play of complex social, cultural, educational, economic and political practices (Balgir 2005). The common beliefs, customs, traditions, values and practices connected with health and disease have been closely associated with the treatment of diseases. In most tribal communities, there is a wealth of folklore associated with health belief.

Tribal populations are particularly vulnerable to malnutrition due to their traditional socio-cultural practices and low literacy level. Several studies on growth and nutritional status were done in rural or urban India (Reddy 2000). Studies on primitive tribes are very few and there is no report on the growth or nutritional status of Bhumia. .

The finding in the present study opens a debatable point about the role of different indices of health and nutritional status assessment. The mean anthropometric measurements indicated that the growth spurt of boys is around 16 years at that age they overcome the girls of same age in both height and weight. Similar trend was reported by various other studies (Hanumanth Rao 1994). The magnitude of wasting in pre-school children was more (36.2%) as compared to NFHS (29.6%) and NNMB (23.7%) report for tribals of Madhya Pradesh and this proportion was observed significantly higher ($P<0.05$) ??NFHS, 1998 and ??NMB, 2000). The wide variation could be due to different sampling techniques used in different studies, similarly the prevalence of chronic energy deficiency was 78% among adults, which was considerably high ($P<0.05$) than the reported figures of 48% among tribals of Madhya Pradesh ??NNMB 2000). The high prevalence of malnutrition observed in the present study could be mainly due to inadequate Dietary intake.

Other reasons could be poor socio-economic status, low purchasing power and faulty feeding habits etc. (Verma 1999). In the present study the intake of cereals was higher than the recommended level. Similar observations were also reported by other authors among tribes of Maharashtra and Bihar (Hanumantha Rao et al. 1992 and Chandrasekhar 1997). This is because most of the tribals diet is a cereal-based diet. Most of the nutrients (calories, protein, Iron etc.) except calcium mean intake were inadequate as compared to RDA. Hanumantha Rao et al. (1993) also reported lower intake of such nutrients in Jenu Kurubas, a primitive tribe of Karnataka. The low value of Carotene and Riboflavin could be due to low intake of green vegetable and negligible amount of milk in their diet. The high calcium value was mainly due to frequent consumption of fetid cassia leaves (Cassia-Tora) by this tribe.

From the above discussion, it can be attributed that the poor growth pattern of the Bumia may be due to the poor socio-economic condition. Most of the Bhumia populations of Madhya Pradesh live without modern health care and transport facilities. Hence, the Bhumia the study area face many health and nutritional hazards due to poverty, illiteracy and ignorance. The health and nutrition status of the Bhumia tribes requires an immediate attention in the implementation of short-term supplementary feeding programmes, general medical, and awareness and health care facilities, improvement of food security are needed to overcome the nutrient deficits. ¹

1

Age group (in Yrs.)	and Sex		
	Male	Female	Total
< 1	1	2	3
1-5	61	49	110
6-10	27	27	54
11-14	29	25	54
15-19	38	31	69
20-24	26	35	61
25-29	45	42	87
30-34	33	37	70
35-39	20	20	40
40-44	16	14	30
45-50	13	12	25
51-60	5	4	9
60+	3	1	4
Total	317	299	616

Figure 1: Table 1 :

2

2013
D D D D) D
(

Age group (in Yrs.)	Male Mean \pm S.D.	Height		Female		Male		Weight		Female Median
		Median	Mean \pm S.D	Median	Mean \pm S.D.	Median	Mean \pm S.D.	Median	Mean \pm S.D	
1+	74.1 \pm 0.2	76.0		76.1 \pm 0.5	76.0	6.2 \pm 1.1	5.8	6.7 \pm 1.3	6.5	
2+	80.4 \pm 4.1	81.5		81.7 \pm 3.4	82.0	9.1 \pm 1.8	8.5	8.8 \pm 1.6	8.8	
3+	85.7 \pm 5.1	85.3		86.5 \pm 7.1	86.0	10.6 \pm 1.5	9.8	10.4 \pm 1.5	9.8	
4+	92.3 \pm 6.6	91.0		93.2 \pm 5.1	93.0	12.0 \pm 2.4	10.5	11.2 \pm 1.8	11.0	
5+	100.5 \pm 7.1	99.0		101.4 \pm 7.0	101.0	14.3 \pm 2.6	11.7	13.1 \pm 2.6	13.0	
6+	102.2 \pm 7.2	101.0		103.2 \pm 6.7	104.0	14.4 \pm 2.1	13.7	13.5 \pm 2.1	14.0	

© 2013 Global Journals Inc. (US)

Figure 2: Table 2 :

3

Indicators	< -3SD	-3SD to<-2SD	-2SD to<-1SD	-1SD to<Median	> Me- dian
	Severe		Moderate	Normal	
Weight for Age (Underweight)	23.2		35.6	28.1	7.6
Height for Age(Stunting)	20.1		22.1	22.6	16.8
Weight for Height (Wasting)	9.0		27.2	27.2	21.3

Figure 3: Table 3 :

4

BMI Grades	Male	Female	Total
CED III (<16.0)	39.6 (100)	35.3 (83)	37.6 (183)
CED II (16.0-17.0)	15.2 (38)	14.7 (35)	15.0 (73)
CED I (17.0-18.5)	23.0(58)	23.6 (55)	23.4 (113)
Low Weight Normal (18.5-20.0)	14.9 (36)	16.4 (38)	15.6 (74)
Normal (20.0-25.0)	6.5 (17)	9.6 (22)	8.0 (39)
Over Weight (>25.0)	0.2 (1)	0.4 (2)	0.3 (3)

Figure 4: Table 4 :

5

Bhumias (gm/cu/day)	Mean \pm SD (n=475)	RDA (ICMR)
Food Items		
Cereals	475.4* \pm 185.5	460
Pulses	28.8* \pm 13.6	40
Green leafy vegetables	29.7* \pm 12.7	40
Roots and Tubers	13.2* \pm 4.6	50
Other vegetables	46.5* \pm 24.4	60
Flesh food	1.7* \pm 1.6	40
Milk and milk products	5.5* \pm 3.2	150
Oil and fats	2.2* \pm 3.6	40
Sugar and jaggery	0.4* \pm 0.1	30
P<0.05		

Figure 5: Table 5 :

6

Nutrient intake	(cu/day) Mean \pm SD (n=475)	RDA (ICMR)
Energy (Kcal)	1810.5* \pm 739.6	2425
Protein (g)	50.4* \pm 21.1	60
Fat (g)	11.1* \pm 6.4	20
Calcium (mg)	435.5 \pm 232.2	400
Iron (mg)	15.7* \pm 13.3	28
Carotene (?g)	368.1* \pm 256.2	2400
Thiamine (mg)	1.3 \pm 0.4	1.2
Riboflavin (mg)	0.4* \pm 0.3	1.4
Vitamin C (mg)	17.8* \pm 13.7	40
P<0.05		

Figure 6: Table 6 :

.1 Acknowledgement

The authors are thankful to all the subjects and their families for their kind cooperation. AJ is grateful to Prof. A. K. Kapoor, Prof. Satwanti Kapoor of Delhi University, Dr A. Jain, and Dr Pradeep Kumar of Sagar, Madhya Pradesh for their help and valuable suggestions during research.

[Weiner and Lourie ()] , J S Weiner , J A Lourie . *Human Biology: A Guide to Field Methods. International Biological Programme, IBP* 1981. Marylebone. (9) .

[Thimmayama ()] *A Handbook of Guidelines in Socio-Economic and Diet Survey*, Bvs Thimmayama . 1987. Hyderabad. National Institute of Nutrition

[Verma (ed.) ()] *A Study of Socio-Cultural Determinants of Population Growth in a Primitive Tribe-The Baiga of Mandla District*, A Verma . Rani Durgawati Vishwa Vidhyalaya (ed.) 1999. Jabalpur. (Ph.D. Thesis)

[Rao et al. ()] 'Assessment of Nutritional Status of Jenu Kurubas -A primitive tribe of Karnataka'. Hanumantha Rao , D Brahmam , Gnv , Mallikharjuna Rao , K , Gal Reddy Ch , Pralhad Rao . *Ind Jr of Nutr Dietet* 1993. 30 p. .

[Jelliffe ()] 'Assessment of the Nutritional Status of the Community'. D B Jelliffe . *Genava: WHO* 1966. (53) .

[Balgir ()] 'Biomedical anthropology in contemporary tribal society of India'. R S Balgir . *Contem-porary Society: Tribal Studies (Tribal Situation in India)*, 2005.

[James et al. ()] 'Definition of Chronic Energy Deficiency in Adults'. Wpt James , Anna Ferro-Luzzi , J C Waterlow . *Eur J Clin Nutr* 1988. 42 p. .

[Elwin ()] V Elwin . *The Baigas. 1st Reprint in*, (India, Delhi) 1986. Gian Publishing House.

[Balgir (ed.) ()] *Genetic markers in the tribal health, development and welfare with special reference to North Eastern India*, R S Balgir . Anthropology, Population and Development. Sharma JC. (ed.) 1995. Ed.) New Delhi: Inter-India Pub. p. .

[Patel ()] *Global Perception of Tribal Research in India*, M L Patel . 2002. New Delhi. p. 37. (Atlantic publisher & Distribution)

[Reddy and Rao ()] 'Growth pattern of the Sugalis a tribal population of Andhra Pradesh'. Pyb Reddy , A P Rao . *India. Ann.Hum. Biol* 2000. 27 p. .

[Health and Nutritional Status of a Primitive Tribe of Madhya Pradesh: Bhumia] *Health and Nutritional Status of a Primitive Tribe of Madhya Pradesh: Bhumia*,

[Health and Nutritional Status of a Primitive Tribe of Madhya Pradesh: Bhumia] *Health and Nutritional Status of a Primitive Tribe of Madhya Pradesh: Bhumia*,

[Srinivasan ()] 'Health care in rural India: Problems and Challenges'. S Srinivasan . *Anthropology Development and Nation Building*, A K Kalla, K S Singh, Eds, Delhi (ed.) 1987. Concept Publishing Company. p. .

[Balgir (ed.) ()] *Human genetics, health and tribal development in Orissa*, R S Balgir . Environment, Health and Development: Anthropological Perspective. Dash Sharma P.Ed. Ranchi (ed.) 2000. p. . SC Roy Institute of Anthropological Studies

[Indian Council of Medical Research (ICMR) 1981. Recommended Dietary Intakes for Indians] *Indian Council of Medical Research (ICMR) 1981. Recommended Dietary Intakes for Indians*, New Delhi: ICMR.

[Indian Council of Medical Research (ICMR) 1990. Recommended Dietary Allowances for Indians-A Report of Expert Group of I] *Indian Council of Medical Research (ICMR) 1990. Recommended Dietary Allowances for Indians-A Report of Expert Group of ICMR*, New Delhi: ICMR.

[Rao and Rao ()] 'Levels of Malnutrition and Socio-Economic Conditions among Maria Gonds'. Hanumantha Rao , D , Mallikharjuna Rao . *J Hum Ecol* 1994. 5 p. .

[Measuring Changes in Nutritional Status. Guidelines for Assessing the Nutritional Impact of Supplementary Feeding Programme] *Measuring Changes in Nutritional Status. Guidelines for Assessing the Nutritional Impact of Supplementary Feeding Programme for Vulnerable Groups*, 1983. Geneva: WHO.

[Mumbai: International Institute of Population Sciences. 19. National Nutrition Monitoring Bureau ()] *Mumbai: International Institute of Population Sciences. 19. National Nutrition Monitoring Bureau*, 1998. 2009. Hyderabad. National Family Health Survey-II India ; National Institute of Nutrition (Diet and Nutritional Status of Tribal Population. Report on first repeat survey)

[Mitra ()] 'National Center for Health Statistics 1976. NCHS Growth Curves for Children 0-18 years'. A Mitra . *No. (PHS) Series Nutrition and Development*. Biswas M & Andersen P (ed.) 1985. DHEW Publication. 17 (165) p. . (The nutrition situation in India)

[Gorden et al. ()] 'Nutrition and infection field study in Guatemalan villages 1959-1964, acute diarrheal diseases and nutrition disorder in general disease incidence'. J E Gorden , W Ascoli , L J Mata , M A Guzman , N S Scrimshaw . *Achieves Environ Hlth* 1968. 16 p. .

5 DISCUSSION

- 195 [Chandrasekhar et al. ()] 'Nutritional status of selected Oraon tribes of Bihar'. U Chandrasekhar , S Kowsalya ,
196 S Sinha . *Ind J Nutri Dietet* 1997. 34 p. .
- 197 [Gopalan et al. ()] *Nutritive Values of Indian Foods*, C Gopalan , B V Ramasastri , S C Balasubramanyam ,
198 Narsinga Rao , B S Deosthale , Y G Panth , KC . 1990. Hyderabad. National Institute of Nutrition (ICMR
- 199 [Rao et al.] 'Prahlad Rao N 1992. Nutritional Status of Maria Gond -A primitive tribe of Maharashtra'.
200 Hanumantha Rao , D Brahmam , Gnv , Mallikharjuna Rao , K , Gal Reddy Ch . *Ind Jr of Nutr Dietet*
201 29 p. .
- 202 [Tewari ()] 'Primitive Tribes of Madhya Pradesh'. D N Tewari . *Ministry of Home Affairs*, (New Delhi; Govt. of
203 India) 1984.
- 204 [Sharma ()] 'The Baiga (the medicine man)'. K Sharma . <http://www.countercurrents.org/sharma261007.htm> *Countercurrents.org* 2007.
- 206 [Balgir ()] 'Tribal health problems, disease burden and ameliorative challenges in tribal communities with special
207 emphasis on tribes of Orissa'. R S Balgir . *Jabalpur: Regional Medical Research Centre for Tribals* 2007. p. .
- 208 [World Health Organization 1990. Diet, Nutrition, and the Prevention of Chronic Diseases WHO Technical Report Series]
209 'World Health Organization 1990. Diet, Nutrition, and the Prevention of Chronic Diseases'. *WHO Technical*
210 *Report Series* (797) .